

## CONTENTS

	Page
Mortality and fertility life tables of leaf folder <i>Cnaphalocrocis medinalis</i> Guenee (Lepidoptera: Pyralidae) in rice: Padmavathi, Gururaj Katti, A. P. Padmakumari I. C. Pasalu. . . . .	1
Records of aphids and aphidocolous ants (Hymenoptera: Formicidae) from Karnataka: Sunil Joshi. . . . .	15
Voltage gated potassium channels in the larval CNS neurons of coconut black headed caterpillar, <i>Opisina arenosella</i> Wlk.: K. Subaharan, Murali Mohan. . . . .	25
Fecundity and developmental stages of Rice Blue Beetle, <i>Leptispa pygmaea</i> Baly (Coleoptera: Chrysomelidae) on rice varieties: K. Karthikeyan, Sosamma Jacob. . . . .	35
Insecticidal efficacy of <i>Chromolaena odorata</i> (compositae) on the coconut beetle, <i>Oryctes rhinoceros</i> (Linn.): S. Leena, B. T. Rayudu, D. Muraleedharan. . . . .	41
Population dynamics, species composition and feeding site preference of rice black bug (hemiptera: pentatomidae) infesting rice: P. Anandhi, M. A. K. Pillai, Savita Varma. . . . .	47
Revival and recharacterization of genus <i>Thyrgorina</i> walker (Lepidoptera: Arctiidae: Arctiinae) and taxonomic studies on four Indian species of this genus from Western Ghats of India: Jagbir Singh Kirti, Navneet Singh Gill. . . . .	53
 SHORT COMMUNICATIONS	
Evaluation of <i>Bacillus thuringiensis</i> Berliner subsp. <i>kurstaki</i> for management of lepidopteran pests of lac insect: A. K. Jaiswal, A. Bhattacharya, S. Kumar, J. P. Singh. . . . .	65
Effect of some selected biopesticides on growth and development of <i>Spodoptera litura</i> fab. (Lepidoptera: Noctuidae) larvae: M. Ravi, N. Dhandapani, N. Sathiah, M. Murugan. . . . .	71
Occurrence of Phthiraptera on the house crow, <i>Corvus splendens</i> (Passeriformes: Corvidae): Sultan Beg, Nidhi Gupta, Sandeep Kumar, Vikram Khan, Shivika Bhatnagar, A K Saxena. . . . .	75
Effect of row spacing and different levels of nitrogenous fertilizer on population density of thrips ( <i>Thrips tabaci</i> Lind.) in onion ( <i>Allium cepa</i> L.): Abhishek Shukla, S. S. Rathore. . . . .	79

Continued on back cover



## ENTOMON

ENTOMON is a quarterly journal of the Association for Advancement of Entomology issued in March, June, September and December, devoted to publication of research work on various aspects of insects and other arthropods.

### EDITORIAL BOARD

C. C. ABRAHAM, Thrissur, India  
T. N. ANANTHAKRISHNAN, Chennai, India  
APARNA DUTTA GUPTA, Hyderabad, India  
K. P. GOPINATHAN, Bangalore, India  
GOVINDAN BHASKARAN, Thiruvananthapuram, India  
KLAUS H. HOFFMANN, Toki, Germany  
G. MADHAVAN NAIR, Thiruvananthapuram, India  
N. MOHANDAS, Thiruvananthapuram, India  
K. S. S. NAIR, Thiruvananthapuram, India  
T. C. NARENDRAN, Calicut, India  
M. S. PALANISWAMI, Thiruvananthapuram, India  
V. K. K. PRABHU, Thiruvananthapuram, India  
K. D. PRATHAPAN, Thiruvananthapuram, India  
R. J. RABINDRA, Bangalore, India  
T. P. RAJENDRAN, New Delhi, India  
V. V. RAMAMURTHY, New Delhi, India  
SEIJI TANAKA, Ibaraki, Japan  
R. V. VARMA, Thiruvananthapuram, India  
**D. MURALEEDHARAN (Managing Editor)**, Thiruvananthapuram, India  
C. A. JAYAPRAKAS (Associate Editor), Thiruvananthapuram, India

Address MS and all editorial correspondence to Managing Editor, ENTOMON, Centre for Arthropod Bioresources and Biotechnology (CABB), University of Kerala, Kariavattom, Trivandrum 695581, India.

### SUBSCRIPTION RATES

Annual subscription for Institutions: Rs. 2000.00 (in India); US\$ 250 (Air Mail)

Annual subscription for individuals: Rs. 300.00 (in India); US\$ 100 (Air Mail)

---

© 2008 by the Association for Advancement of Entomology. All rights reserved

---

1. All remittance to the Journal or Association should be sent to the Secretary-Treasurer of the Association by Bank Draft only. A/c payee in favour of the Association for Advancement of Entomology, payable at Kariavattom.
2. Requests for replacement copies of ENTOMON in lieu of numbers lost in transit, should reach the Secretary-Treasurer not later than three months after the date of publication of the number.

ENTOMON is covered in the following abstracting/indexing journals: *Chemical Abstracts*, *Review of Applied Entomology*, *Science Citation Index* and *Current Contents/Agriculture, Biology and Environmental Sciences*, *Biological Abstracts*, *Entomology Abstracts* and other relevant abstracts, *Referativny Zhurnal* and *Current Advance in Biological Sciences* and *New Entomological Taxa*.



## Mortality and fertility life tables of leaf folder *Cnaphalocrocis medinalis* Guenee (Lepidoptera: Pyralidae) in rice

Padmavathi \*, Gururaj Katti, A. P. Padmakumari and  
I. C. Pasalu

Directorate of Rice Research, Rajendranagar, Hyderabad 500 030, India  
Email: paddi68@rediffmail.com

**ABSTRACT:** Identifying key natural enemies and vulnerable life stages of leaf folder are essential for careful planning and developing practical management programs. The present study is an attempt to construct a life table of natural populations of rice leaf folder in India, to delineate and quantify the role of various mortality factors. Analysis of life table data of rice leaf folder based on natural field populations revealed various mortality factors responsible for the fatality of leaf folder at different stages.  $K$ -values indicated that the total generation mortality was 72.33% in *Dry* and 54.15% in *Wet* seasons, of which biotic factors contributed 34.06% and 29.72% mortality while other factors resulted in 38.27% and 24.43% mortality in the two seasons, respectively. Series of stage-specific mortalities showed maximum mortality in the pupal stage ( $k_{7+8} = 0.2957$  in *Dry* and 0.211 in *Wet* seasons) followed by late larval stage ( $k_{4+5+6} = 0.1716$  in *Dry* and 0.1202 in *Wet* seasons). Bio-mortality factors included seven parasitoids, one pathogen and five predators. Of these, pupal parasitoids viz., *Brachymeria* sp. and *Xanthopimpla flavolineata* Cameron were the key mortality factors causing 34.15% mortality in *Dry* and 32.35% in *Wet* seasons. Survival rates from greenhouse studies indicated higher mortality early in the life cycle i.e., egg and larval stage. Total generation mortality was 26.1% in *Dry* season and 25.9% in *Wet* season indicating that  $\approx 75\%$  of the population entered into the next generation under greenhouse conditions. Intrinsic rate of natural increase ( $r_m$ ) was 0.183 females per female per day and the finite rate of increase ( $\lambda$ ) was 1.2008 females per female per day. © 2008 Association for Advancement of Entomology

**KEYWORDS:** mortality factors, parasitoids, pathogens, net reproductive rate, intrinsic rate of increase, survival rates, *Cnaphalocrocis medinalis*

### INTRODUCTION

Rice leaf folder, *Cnaphalocrocis medinalis* Guenee is widely distributed in all the rice growing tracts of South and South-east Asia. Once considered as minor pest,

\*Corresponding author

it has now attained a major pest status due to intensification of rice cultivation and outbreaks of this pest have also been reported (Khan *et al.*, 1988; Senapati and Panda, 2006). Life tables provide an important tool in understanding the changes in population of insect pests during different developmental stages throughout their life cycle. Identifying key natural enemies and vulnerable life stages of leaf folder are essential for possible biological control efforts and for careful planning and developing practical management programs (Midega *et al.*, 2005; Medieros *et al.*, 2000; Hu *et al.*, 2000). Characterization of stage specific survivorship is also important for reflecting the impact of natural enemies in economic thresholds (Pedigo, 1997).

Determination of population increase from reproductive capacity is another crucial component in the study of insect populations. Population increase can be described by a fertility table that gives the potential reproductive ability of females at different times. A standard estimator for the growth rate of insect populations is the intrinsic rate of increase ( $r_m$ ), which describes the maximal rate of increase at any time interval under optimal conditions. The rate of multiplication of a population per generation is described by the net reproductive rate ( $R_o$ ), which is dependent on  $r_m$  (Southwood, 1978).

Though there are a few earlier reports on biology and life cycle aspects of leaf folder (Barrion *et al.*, 1991; Pang *et al.*, 1981), information on life table of the pest is lacking. The present study was carried out to determine the variables of life and fertility tables to be used as a basis for understanding the population dynamics of *C. medinalis* in rice and may help in finding the reasons for flare up of this pest.

## MATERIALS AND METHODS

### Mortality tables

Greenhouse and field studies were conducted at Directorate of Rice Research, Hyderabad during Wet (*Kharif*) and Dry (*Rabi*) seasons of the years, 2004 and 2005.

### Greenhouse studies

To assess the inherent mortality of rice leaf folder, age-specific life-tables were constructed from the data recorded on the same cohort under the greenhouse conditions ( $23 \pm 10^\circ\text{C}$  temperature and  $60 \pm 10\%$  relative humidity) coinciding with Wet and Dry seasons. Paired adults were released for oviposition on 20–25 day old susceptible variety i.e., Taichung Native-1 (TN 1) plants placed in wooden cages (66 cm length  $\times$  97 cm width  $\times$  86 cm height) having wire mesh on all sides. Eggs from the same cohort were considered for this life table study. There were three replications in each season. The leaves of plants with eggs were tagged and these potted plants were kept in wooden cages in the greenhouse. The tagged eggs were observed daily till they hatched. The larvae were observed till they pupated. Mortality data were recorded at each stage. The pupae were collected and placed in test tubes and were observed for the emergence of adult moths. The number of males and females emerged were counted to know the sex ratio.

### *Field studies*

The experimental field consisted of 750 m<sup>2</sup> area with a susceptible variety i.e., Taichung Native-1 (TN 1) planted in nine blocks of 80 m<sup>2</sup> area each. Each block is considered as one replication and thus there were nine replications per season. The rice crop was raised following recommended agronomic practices without any plant protection measures. Absolute populations of leaf folder and that of its natural enemies were recorded in both Wet and Dry seasons of both the years. Due to overlapping generations of the pest, life tables were prepared for seasons instead of generations. In each block, one square meter area was marked at two places, of which one was kept open and the other area was covered on all four sides by a polyethylene sheet to restrict the larval movement. The top portion was kept open for movement of natural enemies. The stages sampled in the field included different larval instars, pupae and adults that emerged from collected pupae. Observations were initiated at first instar stage and continued till pupation, at four-day interval. All the leaves of each plant (40 hills) in one square meter area were inspected for folded leaves with larvae. The number of healthy surviving larvae in folded leaves was counted and allowed to remain in the field, while the morbid/parasitised larvae were collected and reared individually in laboratory, for emergence of adult parasitoids. Similarly, in subsequent observations the healthy larvae that pupated were collected and brought to laboratory and observed for adult emergence while parasitized pupae were separated and pupal parasitism was estimated. The number of males and females that emerged from pupae was counted to determine the sex ratio. The emerged adults were then paired and kept for oviposition.

To assess the mortality at egg stage, pots with TN 1 plants having leaf folder eggs (0–24 h old) were placed in the field at random for about two days. Predators found during sampling were also visually counted to provide an estimate of potential predators in the study area. At larval stage, the differences in larval population observed in open and closed areas were considered due to larval dispersal. Based on these data, stage specific mortalities were worked out after grouping first two instars of *C. medinalis* larvae as 'early instars' and rest three as 'late instars'. Analysis to recognize the role of various factors in life table was done as suggested by Varley and Gradwell (1960). Series of stage specific mortalities ( $k$ -values) for egg, larvae and pupae were then summed up to compute the total generation mortality ( $K$ ) in the life-table of the leaf folder. Means were computed by pooling the data of four seasons of both the years.

To know the effect of abiotic factors on leaf folder population, meteorological data viz., maximum temperature, minimum temperature, relative humidity (morning and evening), rainfall and sunshine hours were recorded. Correlations were worked out between larval population and the weather parameters.

### **Fertility table**

Age specific fertility or fecundity parameters were recorded on TN 1 variety under greenhouse conditions. Leaf folder adults that emerged on the same day from the field collected pupae were paired and released for oviposition on 20–25 day old TN 1 plants

covered with cylindrical mylar cage. Adults were fed with 20% honey solution inside cages. Ten such pairs were used for the study in each season. The number of eggs laid by a female was counted and adults were released daily on fresh plants till their death. After hatching, larvae were allowed to develop by providing fresh TN1 plants as and when necessary. The number of females born (mx) out of the eggs laid by a female was calculated based on these observations.

Based on the data obtained, mortality and fertility life-tables were prepared according to Harcourt (1969); Southwood (1978); Hu *et al.* (2000); Medieros *et al.* (2000) and Midega *et al.* (2005). The net reproductive rate ( $R_o$ ), approximate generation time ( $T_c$ ) and capacity for increase ( $r_c$ ) were calculated as per Ghosh *et al.* (1995) and Mohanty *et al.* (1995). The accurate value of  $r_m$  was calculated by taking two arbitrary trial values on either side of the value for  $R_o$  differing only in the second place and substituting them in the formula. Graphically  $r_m$  value was estimated according to Narendra Reddy and Singh (1998). By using accurate  $r_m$  value, the true generation time ( $T$ ) and finite rate of increase ( $\lambda$ ) were further calculated. The time taken (in days) for the leaf folder population to double was also worked out.

## RESULTS

### Mortality tables

#### *Greenhouse studies*

Life-table of leaf folder under greenhouse conditions is summarized in Table 1. Females lived for about a week with pre-oviposition period of 3–4 days and oviposition period of 4–5 days. Total development period from egg to adult was found to be 24.8 days. Survival rates indicated higher mortality early in the life cycle i.e., egg and larval stage (Fig. 1). Hatching percentage of 84.69 and 82.03 was found in the two seasons. Total generation mortality was 26.1% in *Dry* season and 25.9% in *Wet* season indicating that *ca.* 75% of the population entered into the next generation.

#### *Field studies*

Analysis of field life table data based on natural populations revealed several mortality factors responsible for the fatality of leaf folder at different stages (Table 2). K-values indicated that the total generation mortality was 72.34% in *Dry* and 54.15% in *Wet* seasons, of which biotic factors contributed 34.06% and 29.72% mortality while other factors resulted in 38.28% and 24.43% mortality in the two seasons, respectively. Series of stage-specific mortalities showed maximum mortality in the pupal stage ( $k_{7+8} = 0.2957$  in *Dry* season and 0.211 in *Wet* season) followed by late larval stage ( $k_{4+5+6} = 0.1716$  in *Dry* and 0.1202 in *Wet* seasons).

Bio-mortality factors included seven parasitoids, one pathogen and five predators. Of these, pupal parasitoids *viz.*, *Brachymeria* sp., *Xanthopimpla flavolineata* Cameron and *Tetrastichus* sp. were the key mortality factors causing 34.05% mortality in *Dry* and 32.85% in *Wet* seasons of the total pupal mortality (Fig. 2). Parasitism was

TABLE 1. Life table of leaf folder under greenhouse conditions

Stage of the insect (x)	*Number living (lx) Mean $\pm$ SE(m)		Factors responsible for dx (dxf)	*Number dying (dx)		dx as % of lx (100 qx)		kappa (k) values	
	Dry season <sup>\$</sup>	Wet season <sup>#</sup>		Dry season	Wet season	Dry season	Wet season	Dry season	Wet season
Eggs	122 $\pm$ 1.15	85.34 $\pm$ 0.33	unhatched	18.67 $\pm$ 0.34	15.34 $\pm$ 0.33	15.30	17.97	0.073 (k <sub>1</sub> )	0.086 (k <sub>1</sub> )
Larvae	103.34 $\pm$ 1.76	70 $\pm$ 2.88	unknown/natural	19.34 $\pm$ 0.88	11.67 $\pm$ 0.34	18.70	16.67	0.090 (k <sub>2</sub> )	0.0792 (k <sub>2</sub> )
Pupae	84 $\pm$ 1.15	58.34 $\pm$ 0.88	unknown/natural	11.67 $\pm$ 0.34	8 $\pm$ 0.57	13.88	13.71	0.065 (k <sub>3</sub> )	0.0641 (k <sub>3</sub> )
Adults	72.34 $\pm$ 1.76	50.34 $\pm$ 0.34	deformed	5.34 $\pm$ 0.33	3.34 $\pm$ 0.33	7.37	6.62	0.033 (k <sub>4</sub> )	0.0297 (k <sub>4</sub> )
Adults	67 $\pm$ 1.16	47 $\pm$ 0.57							
				Total generation mortality (K)				0.261	0.259

\*Mean of two years (three replications/season)

<sup>\$</sup> Dry season (*Rabi*) – November to April of 2004 and 2005<sup>#</sup> Wet season (*Kharif*) – July to October of 2004 and 2005




FIGURE 1. (a) Survivorship curve – greenhouse; (b) Survivorship curve – Field.

also high in late larval instars (20.88% in *Dry* and 17.27% in *Wet seasons*) due to *Macrocentrus philippinensis* Ashmead and *Goniozus* sp. In early instars, *Apanteles angustibasis* (Gahan) and *Apanteles* sp. caused 8.54% (*Dry season*) and 8.7% (*Wet season*) mortality. The lone pathogen recorded was *Beauveria bassiana* (Balsamo) Vuillemin causing 3.71% and 1.89% mortality of late larval instars in the *dry* and *wet* seasons, respectively. Predators included one spider (*Tetragnatha* sp.), beetles (*Paederus* sp., *Ophionea* sp.) and damsel flies at all the stages.

Larval dispersal was relatively higher during early instars (8.24 to 12.93%) compared to late instars (6.59 to 11.57%) in both the seasons. Blackening of pupae resulted in the mortality of pupae (23.16% in *Dry* and 9.08% in *Wet*) while



TABLE 2. Life table of rice leaf folder under field conditions

Stage of the insect (x)	*Number living (lx)		Factors responsible for dx (dx/f)	*Number dying (dx)		dx as % of lx (100 qx)		Kappa (k) values	
	Dry season <sup>\$</sup>	Wet season <sup>#</sup>		Dry season	Wet season	Dry season	Wet season	Dry season	Wet season
Eggs	109.33 ± 0.67	106.12 ± 0.45	Unhatched	23.33 ± 0.57	20.25 ± 0.59	21.34	19.08	0.1042(k <sub>1</sub> )	0.0920(k <sub>1</sub> )
Early instar larvae	86 ± 0.62	85.87 ± 0.71	Parasitism	7.77 ± 0.32	7 ± 0.46	9.04	8.15	0.0411(k <sub>2</sub> )	0.0369(k <sub>2</sub> )
	78.23 ± 0.52	78.87 ± 0.56	Dispersal	10.11 ± 0.42	6.5 ± 0.33	12.93	8.24	0.0602(k <sub>3</sub> )	0.0374(k <sub>3</sub> )
Late instar larvae	68.11 ± 0.61	72.37 ± 0.44	Parasitism	14.22 ± 0.32	12.5 ± 0.37	20.88	17.27	0.1018(k <sub>4</sub> )	0.0823(k <sub>4</sub> )
	53.88 ± 0.63	59.87 ± 0.59	Pathogen	2 ± 0.37	1.13 ± 0.29	3.71	1.89	0.0164(k <sub>5</sub> )	0.0083(k <sub>5</sub> )
	51.88 ± 0.54	58.74 ± 0.53	Dispersal	6 ± 0.62	3.87 ± 0.29	11.57	6.59	0.0534(k <sub>6</sub> )	0.0296(k <sub>6</sub> )
Pupae	45.88 ± 0.48	54.87 ± 0.59	Parasitism	15.67 ± 0.53	17.75 ± 0.36	34.15	32.35	0.1813(k <sub>7</sub> )	0.1697(k <sub>7</sub> )
	30.22 ± 0.40	37.12 ± 0.56	Blackening	7 ± 0.41	3.37 ± 0.32	23.16	9.08	0.1144(k <sub>8</sub> )	0.0413(k <sub>8</sub> )
Adults	23.22 ± 0.40	33.75 ± 0.44	Deformation	2.55 ± 0.26	3.25 ± 0.25	10.98	9.63	0.0505(k <sub>9</sub> )	0.0440(k <sub>9</sub> )
	20.67 ± 0.50	30.50 ± 0.56							
Total generation mortality (K)						0.7234	0.5415		

\*Mean of two years (nine replications/season)

<sup>\$</sup> Dry season (Rabi) – November to April of 2004 and 2005<sup>#</sup> Wet season (Kharif) – July to October of 2004 and 2005




FIGURE 2. Stage specific parasitism of rice leaf folder.

deformation of adults was another factor accounting for 10.98% in Dry season and 9.63% mortality in Wet season.

Correlation between leaf folder population and weather factors indicated that the maximum temperature ( $r = -0.6977$ ), minimum temperature ( $r = -0.5781$ ) rainfall ( $r = -0.6702$ ) and sunshine hours ( $r = -0.5798$ ) have significant negative relationship while morning relative humidity ( $r = 0.6719$ ) and evening relative humidity ( $r = 0.7501$ ) exhibited positive relationship.

### Fertility table

Female moth started laying eggs after 27th day of pivotal age which lasted up to 33rd day with lx values being 0.56 to 0.04, respectively (Table 3). The maximum mean progeny production per day (Vx) was 71.93 females per female on 29th day of pivotal age. The net reproductive rate ( $R_0$ ) was 210.77 indicating that the population of *C. medinalis* would be able to multiply 210.77 times per generation (Table 4). The mean length of generation ( $T_c$ ) was 29.24 days. Intrinsic rate of natural increase ( $r_m$ ) was derived as 0.183 females per female per day (Table 4 and Fig. 3). The finite rate of increase ( $\lambda$ ) was 1.2008 females per female per day. The population multiplied 3.6 times per week and doubled in 4.75 days.

TABLE 3. Age specific fecundity table of *Cnaphalocrocis medinalis*

Pivotal age in days (x)	$I_x$	$m_x$	$V_x$			Trial $r_m$ values		Accurate value
			$(l_x m_x)$	$x.l_x m_x$	$x$	$r_m = 0.18$	$r_m = 0.19$	
						$e^{7-r} m^x . l_x m_x$	$e^{7-r} m^x . l_x m_x$	
0-25	0.56	—	—	—	—	—	—	—
26	0.56	—	—	—	—	—	—	—
27	0.56	—	—	—	—	—	—	—
28	0.56	94.06	52.67	1474.76	373.92	282.60	343.7952	
29	0.56	128.45	71.93	2085.97	426.53	319.16	390.9945	
30	0.42	141.82	59.56	1786.80	295.00	218.54	269.6120	
31	0.37	63.05	23.33	723.23	96.52	70.79	87.9475	
32	0.12	24.02	2.88	92.16	9.95	7.23	9.0412	
33	0.04	10.15	0.40	13.20	1.15	0.83	1.0457	
Total			210.77	6176.12	1203.07	899.15	1102.4361	

 $l_x$ —Number of living females $m_x$ —Number of living females born per female in each age interval $V_x$ —Total number of female births in each age interval $r_m$ —Intrinsic rate of increase

TABLE 4. Life table parameters

S.No	Population growth statistics	Formula	Calculated value
1	Net reproductive rate ( $R_0$ )	$\Sigma l_x m_x$	210.77
2	Mean length of generation ( $T_c$ )	$\Sigma x l_x m_x / R_0$	29.30 days
3	Capacity for increase ( $r_c$ )	$\text{Loge } R_0 / T_c$	0.1826 females/ female/ day
4	Arbitrary $r_m$ values	0.18 & 0.19	
5	Intrinsic rate of increase ( $r_m$ )	$\Sigma e^{7-r} m^x . l_x m_x$	0.1830 females/ female/ day
6	Finite rate of increase ( $\lambda$ )	$\text{Antilog } e^{r_m}$	1.20081
7	Weekly multiplication rate	$(\lambda)^7$	3.60014
8	Corrected generation time ( $T$ )	$\text{Log } e R_0 / r_m$	29.24 days
9	Doubling time	$\text{Log } e^2 / r_m$	4.7463 days

## DISCUSSION

The present study is an attempt to construct a life table of natural populations of rice leaf folder in India, to delineate and quantify the role of various mortality factors. Greenhouse studies following the fate of a cohort from birth to death revealed that only 75 per cent of the leaf folder population entered into the next generation (Table 1). There was no difference in the mortality across the seasons under greenhouse conditions. In a similar greenhouse study, Chitra *et al.* (2003) also reported that in *C.*




FIGURE 3. Determination of intrinsic rate of increase ( $r_m$ ) of rice leaf folder.

*medinalis*, only 73 per cent reached adulthood whereas 65 per cent reached adulthood in *M. patnalis*. Natural population data under field conditions revealed that a number of biotic and other factors were found causing mortality of rice leaf folder at different stages. Total generation mortality ( $K$ ) was high in *Dry* season as compared to *Wet* season (Table 2) and 28% of population in *Dry* and 48% in *Wet* survived. Mortality due to biotic factors was more in *Dry* season (34.06%) as compared to *Wet* season (29.72%). Pupal and late larval parasitoids viz., *Brachymeria* sp., *X. flavolineata* and *M. philippinensis* were the key mortality factors in both the seasons in both the years (Fig. 2). These findings were in corroboration with Pathummal Beevi *et al.* (2001) who reported 15 to 44% pupal parasitism by *Brachymeria tibialis*, *B. excarinata* and *Tetrastichus* sp and 27.2% larval parasitism by *Cardiochiles philippinensis*. Manisegaran *et al.* (1997) recorded natural parasitism of leaf folder and reported 12.81% parasitism by *A. angustibasis*, 14.87% by *Brachymeria* sp., 15.42% by *Elasmus johnstoni* and 22.42% by *Goniozus* sp. Similarly Kraker *et al.* (1999b) recorded 14 to 56% larval parasitism of leaf folders in irrigated rice in Philippines and the braconid *M. philippinensis* was the most commonly found larval parasitoid. Matsumara (1992) reported larval parasitism by *Apanteles baoris* and *Pediobius mitsukurii* as important mortality factor in the life tables of the migrant skipper, *Parnara guttata guttata* in rice. Parasitisation of various stages and failure of neonate larvae to enter the midribs were the key mortality factors reported in the life tables of sugarcane top borer, *Scirpophaga excerptalis* (Mukunthan, 1989). Similarly, life table studies on the soybean leaf folder, *Nacoleia* sp. revealed that maximum mortality (17–18%) was due to pupal parasitoids (Thakur and Gangwar, 1989).

Survivorship curves for *C. medinalis* indicated higher mortality in early stages of the life cycle under greenhouse conditions, designated as Type IV category (Slobodkin, 1962) whereas mortality was high in late stages of the life cycle under field conditions indicating Type-I category (Fig. 1).

In the present study there was no mortality in egg stage due to parasitism or predation but non-hatching of eggs contributed up to 15.3 to 17.97% mortality. However, Kraker *et al.* (1999a) reported that the egg disappearance was the major mortality factor and the average disappearance of exposed laboratory laid eggs was 40% while that of field-laid eggs was 46%. Egg mortality due to parasitism ranged between 15–18% whereas non-hatching of eggs was of minor importance. The white muscardine fungus, *Beauveria bassiana* was found causing late larval mortality in the present study (Fig. 2). Similarly, Ambethgar (1997) recorded 13–100% mycoses in larval populations due to white muscardine fungus.

Predators recorded include *Paederus* sp. (staphylinid), *Ophionea* sp. (carabid), and, *Tetragnatha* sp (spider). Earlier Rai *et al.* (2000) reported *Paederus fuscipes* as an active predator with a feeding potential of 0.6 rice leaf folder larvae per individual per day. Pathummal Beevi *et al.* (2001) reported that out of 35 predators recorded *Ophionea nigrofasciata*, *Nabis capsiformis* and *Paederus fuscipes* are important. The net reproductive rate ( $R_o$ ) was 210.77. The approximate generation time ( $T_c$ ) was 29.3 days leading to ' $r_c$ ' value of 0.1826 which revealed the capacity for increase of females/female/day. The intrinsic rate of increase  $r_m$  was derived as 0.1830 (Table 4 and Fig. 3). The capacity for increase was slightly less than intrinsic rate of increase indicating population tending towards overlapping generations.

A significant negative correlation was observed between maximum temperature, minimum temperature, rainfall, sunshine hours and leaf folder larval population whereas positive relationship was found between relative humidity (morning and evening) and larval population. Similar relationships were observed by Balasubramanian and Balasubramanian (1985) between leaf folder damage and weather factors. They also indicated that increase in maximum temperature by 1 °C may reduce the leaf folder damage by 1.75 with  $R^2$  value of 0.445.

The life table data, presented here, revealed that the population of rice leaf folder is governed by a number of factors. Mortality was maximum in dry season as compared to wet season. Highest mortality was due to pupal and larval parasitism. Other mortality factors include blackening of pupae due to unknown reason, larval dispersal and diseased larvae. The short doubling time and high intrinsic rate of increase indicated that the leaf folder has a potential to become a serious pest but for the various biotic factors that regulate the population. But if the fragile factors are tinkered of insecticide intervention, the pest may flare up. Thus it is evident that biotic factors play a major role in the population dynamics and conservation of late larval and pupal parasitoids in the field is crucial for efficient management of this pest in rice.

## ACKNOWLEDGEMENTS

We would like to thank the Project Director, Directorate of Rice Research, Hyderabad for providing necessary facilities and Dr. Debjani Dey, Senior Scientist, Division of Entomology, Indian Agricultural Research Institute, New Delhi for the identification of parasitoids.

## REFERENCES

- Ambethgar, V. (1997) Record of white muscardine fungus, *Beauveria bassiana* (Bals.) Vuill. on rice leaf folder complex from Karaikal, Pondicherry Union Territory (India). *Journal of Entomological Research* **21**: 197–199.
- Balasubramanian, G. and Balasubramanian, M. (1985) Surveillance of major pests of rice and cotton. In: *Integrated Pest and Disease Management*, Jayaraj, S. (Ed). Tamilnadu Agricultural University, 363–366.
- Barrior, A.J., Litsinger, J.A., Medina, E.B., Aguda, R.M., Bandong, J.P., Pantua, P.C., Viajante, V.D., Cruz, C.G., Vega, C.M., Soriano, J.S., Camanag, E.E., Saxena, R.C., Tryon, E.H. and Shephard, B.M. (1991) The rice leaf folders, *Cnaphalocrocis medinalis* and *Marasmia* sp (Lepidoptera: Pyralidae) complex in the Philippines: Taxonomy, bionomics and control. *Philippine Entomologist* **8**: 987–1074.
- Chitra, N., Soundararajan, R.P. and Gunathilagaraj, K. (2003) Life table studies for rice leaf folders. *Madras Agricultural Journal* **90**: 371–373.
- Ghosh, S.K., Rao, J., Prakash, A. and Sinha, R.K. (1995) Age specific fecundity, life tables and intrinsic rate of increase of rice tarsonemid mite, *Tarsonemus cuttacki* Iswari (Acari: Tarsonemidae) under greenhouse conditions. *Oryza* **32**: 98–100.
- Harcourt, D.G. (1969) The development and use of life tables in the study of natural insect populations. *Annual Review of Entomology* **14**: 175–196.
- Hu, X.N., Liang, G.W. and Pang, X.F. (2000) Studies on the continuous natural population life tables of Asian corn borer. *Acta Ecologica Sinica* **20**: 107–109.
- Khan, Z.R., Barrior, A.T., Litsinger, J.A., Castilla, N.P. and Joshi, R.C. (1988) A bibliography of rice leaf folders (Lepidoptera: Pyralidae). *Insect Science and its Application* **9**: 129–174.
- Kraker, J.de., Huis, Avan., Lenteren, J.C.van., Heong, K.L. and Rabbinge, R. (1999a) Egg mortality of rice leaf folders *Cnaphalocrocis medinalis* and *Marasmia patnalis* in irrigated rice fields. *Biocontrol* **44**: 449–471.
- Kraker, J.de., Huis, Avan., Heong, K.L., Lenteren, J.C.van. and Rabbinge, R. (1999b) Population dynamics of rice leaf folders (Lepidoptera: Pyralidae) and their natural enemies in irrigated rice in the Philippines. *Bulletin of Entomological Research* **89**: 411–421.
- Manisegaran, S., Letchoumanane, S. and Mohamed Hanifa, A. (1997) Natural parasitism of rice leaf folder, *Cnaphalocrocis medinalis* (Guenee) in Karaikal region. *Journal of Biological Control* **11**: 73–75.
- Matsumara, M. (1992) Life tables of migrant skipper *Parnara guttata guttata* Bremer et Grey (Lepidoptera: Hesperidae) in the northern peripheral area of its distribution. *Applied Entomology and Zoology* **27**: 331–340.
- Medeiros, R.S., Ramalho, F.S., Lemos, W.P. and Zanuncio, J.C. (2000) Age-dependent fecundity and life-fertility tables for *Podisus nigrispinus* (Dallas) (Heteroptera: Pentatomidae). *Journal of Applied Entomology* **124**: 319–324.
- Midega, C.A.O., Ogot, C.K.P.O. and Overholt, W.A. (2005) Life tables, key factor analysis and density relations of natural populations of the spotted maize stem borer, *Chilo partellus* (Lepidoptera: Crambidae) under different cropping systems at the Kenyan coast. *International Journal of Tropical Insect Science* **25**: 86–95.

- Mohanty, B.N., Patnaik, N.C. and Senapati, B. (1995) Life-Budget analysis of the rice hairy caterpillar, *Nisaga simplex* Walker (Lepidoptera: Eupterotidae) in Kalahandi District, Orissa (India). *Journal of Entomological Research* **19**: 1–7.
- Mukunthan, N. (1989) Life tables of sugarcane top borer *Scirpophaga excerptalis* Walker. *Insect Science and its Application* **10**: 269–276.
- Narendra Reddy, C. and Singh, T.V.K. (1998) Rate of increase and stable-age distribution of *Plutella xylostella* on cabbage at Hyderabad. *Indian Journal of Entomology* **60**: 329–333.
- Pang, X.F., Hou, R.H., Liang, G.W. and Li, Z.H. (1981) Studies on the population dynamics and control of rice leaf roller in Hailing island, Guangdong Province. Life table data and its component analysis. *Journal of South China Agricultural College* **2**: 71–84.
- Pathummal Beevi, S., Hazarika, L.K. and Katti, G.S. (2001) Augmentative biocontrol within paddy IPM-Indian Scenario. In: *Augmentative Biocontrol*, Singh, S. P., Murphy, S. T. and Ballal, C. R. (Eds). CABI Bioscience, UK and PDBC, India, 93–117.
- Pedigo, L.P. (1997) General models for economic thresholds. In: *Economic thresholds for integrated pest management*, Higley, L. G. and Pedigo, L. P. (Eds). University of Nebraska Press, Lincoln, NE, 41–57.
- Rai, M.K., Ramamurthy, V.V., Gupta, S.L. and Dey, D. (2000) Preliminary investigations on *Paederus fuscipes* (Curtis) a predator in rice agroecosystems. *Annals of Plant Protection Sciences* **8**: 8–12.
- Senapati, B. and Panda, S.K. (2006) Insect pest management in rice. *Journal of Rice Research* **1**: 72–82.
- Slobodkin, L.B. (1962) *Population and community ecology*. Holt, Rinehart and Winston, New York.
- Southwood, T.R.E. (1978) *Ecological methods with particular reference to the study of insect populations*, 2nd edition, Chapman and Hall, London, 524.
- Thakur, N.S.A. and Gangwar, S.K. (1989) Life table studies on the leaf folders *Nacoleia* spp in soybean in khasi hills of Meghalaya. *Indian Journal of Plant Protection* **17**: 277–278.
- Varley, G.C. and Gradwell, G.R. (1960) Key factors in population studies. *Journal of Animal Ecology* **29**: 399–401.

(Received 26 November 2007; accepted 15 February 2008)







## Records of aphids and aphidocolous ants (Hymenoptera: Formicidae) from Karnataka

Sunil Joshi

*Project Directorate of Biological Control, Post Box No 2491, H.A. Farm Post,  
Bellary Road, Bangalore 560 024, India  
Email: sunjoshi\_pdbc@rediffmail.com*

**ABSTRACT:** A total 66 aphid species under 38 genera belonging to seven subfamilies were recorded in Karnataka in an intensive survey for aphids during 2000–2003. The subfamily Chaitophorinae was recorded for the first time from south India. Eighteen species are recorded for the first time from Karnataka, of which six species are new records from south India. Eleven species of ants belonging to eight genera were found associated with 24 species of aphids infesting 42 host plants. The ants belonged to the subfamilies Formicinae (five species), Myrmicinae (four species) and Dolichoderinae (two species). The genera *Camponotus*, *Crematogaster* and *Monomorium* were more abundant with two species under each genus. *Camponotus compressus* Fabricius was the most common species, attending 15 species of aphids. This was followed by *Monomorium* sp. and *Solenopsis geminata* Fabricius, which attended nine and eight species of aphids, respectively. *Camponotus rufoglaucus* Jerdon, *Crematogaster soror* Forel and *Oecophylla smaragdina* Smith attended three aphid species each. This study records 26 new aphid and ant associations.  
© 2008 Association for Advancement of Entomology

**KEYWORDS:** Aphids, host plants, aphidocolous ants

### INTRODUCTION

In south India, aphid studies started in 1930s and were continued up to 1974, when several species were recorded from Tamil Nadu and a few from Karnataka (Ghosh, 1986). Earlier the aphid fauna of Karnataka was studied by Krishnamurti (1928, 1930, 1950) and Gadiyappanavar (1970). Numerous surveys, across the different seasons were conducted during 2000–2003 in various regions of the state. The areas surveyed for aphids includes rural and urban Bangalore, Belgaum, Chikmagalur, Dakshin Kannada, Dharwad, Hassan, Kolar, Kodagu, Mandya, Mysore and Tumkur.

The presence of ants in the aphid colony helps to increase the aphid population in three different ways. While attending the aphids for honeydew, the ants directly accelerate the aphid's growth and reproduction rate (El-Ziady, 1960). The ants

indirectly increase the reproduction rate by prolonging the aphids stay on young plant tissue where they reproduce faster than older tissues (Banks, 1958). The ants drive away the aphid natural enemies and thus allow the aphid population to increase (Flanders, 1951).

The ant-aphid association has attracted greater attention from the European workers, with the result, most of the literature available on this subject refers to the European aphid species. Various ecological (Curtright, 1965; Skinner and Whittaker, 1981), physiological (Broadbent, 1951) and ethological (Banks, 1958; Banks and Nixon, 1958; Way, 1963; Banks and Macaulay, 1967; Addicott, 1978) studies on aphid ant mutualism have been made from European countries.

Accounts of aphidicolous ants are available from Orissa (Roy and Behura, 1980), Assam, Meghalaya, Rajasthan (Kurl and Misra, 1980), West Bengal, Sikkim (Datta *et al.*, 1982, 1983; Datta and Raychaudhuri, 1984, 1985), Manipur (Devi and Singh, 1986a,b; Devi *et al.*, 1987; Shantibala *et al.*, 1995; Devi *et al.*, 2001), Jammu and Kashmir (Matta and Bhagat, 2000) and Uttranchal (Bisht *et al.*, 2002).

In southern parts of India Gadiyappanavar (1970) and Musthak Ali and Sharatchandra (1986) studied the role of establishment of root aphid on finger millet through ant association. Veeresh and Musthak Ali (1990) gave a list of aphidicolous ants from south India. In Bangalore, Karnataka, Verghese and Tandon (1987) studied interspecific association of *Aphis gossypii* Glover, *Cheilomenes sexmaculata* (Fabricius) and *Camponotus comprassus* Fabricius in guava ecosystem. No attempt has so far been made from south India to study fauna of aphidicolous ants and hence efforts were made to collect and identify aphidicolous ants from Karnataka.

## MATERIALS AND METHODS

### Collection and preservation of aphids

Regular surveys were made to collect aphids and associated ants on various host plants including cultivated crops, ornamental plants, trees and weeds in different districts of Karnataka State viz., Bangalore Rural, Bangalore Urban, Belgaum, Chikmagalur, Dakshin Kannada, Dharwad, Hassan, Kolar, Kodagu, Mandya, Mysore, Tumkur, Bidar and Gulbarga during 2000–2003. Aphids were collected either directly from the plants with the help of camel hair brush or the alate aphids were trapped in yellow pan water traps. Sometimes, the infested plant parts along with aphid colony were collected in small plastic containers or polythene bags. The aphids thus collected were preserved in 70% ethanol in tightly stoppered small plastic vials. For microscopic studies the aphids were processed following the method proposed by van Emden (1972).

### Systematics and identification of aphid species

The aphids recorded in the present study were grouped in one of the nine subfamilies Aphididae viz., Eriosomatinae (Pemphiginae), Hormaphidinae, Anoeciinae, Chaitophorinae, Drepanosiphinae, Greenideinae, Aphidinae, Phloeomyzinae and

Lachninae. The classification proposed by Remaudière and Remaudière (1977) with subfamily names used by Nieto Nafria *et al.* (1998) are followed in the study.

The aphids were identified with the help of keys provided by Eastop (1966), Raychaudhuri (1980) and Martin (1983).

The infested plant parts along with aphid colony and associated ants were collected in small plastic containers (12 × 7 cm) or polyethylene bags. The plant parts were kept in an acrylic sheet cage (30 × 30 × 30 cm) after bringing them to the laboratory. The ants were separated from aphid colony and preserved in 70% ethyl alcohol in small plastic vials (2.5 ml). Each vial was furnished with data on locality, host plant, date and collector's name on thin paper.

## RESULTS AND DISCUSSION

A total of 66 aphid species under 38 genera belonging to seven subfamilies were recorded from Karnataka in the present study (Table 1). Krishnamurti (1928, 1930, 1950) and Gadiyappanavar (1970) recorded 20 and 30 aphid species from Karnataka, respectively. Gadiyappanavar (1970) also provided a list of 67 host plants of various aphid species.

The subfamily Chaitophorinae was recorded for the first time from south India, however, the subfamilies Phloemyzinae and Anoeciinae could not be recorded. In earlier studies also species belonging to these subfamilies were not recorded. Family Aphidinae was the most dominant group with 65.15% of the total species recorded. Subfamily Greenideinae and Hormophidinae were second and third in dominance, respectively, having 12.12 and 9.09% of the total species recorded.

Out of 66 aphid species recorded in the present study, 24 species were attended by 11 species of ants belonging to eight genera on 42 species of host plants (Table 2). The ants belonged to the subfamilies Formicinae (five species), Myrmicinae (four species) and Dolichoderinae (two species). The genera *Camponotus*, *Crematogaster* and *Monomorium* were more common with two species each. *Camponotus compressus* Fabricius was the most common, attending 16 species of aphids. This was followed by *Monomorium* sp. and *Solenopsis geminata* Fabricius, which attended nine and eight species of aphids, respectively. *Camponotus rufoglaucus* Jerdon, *Crematogaster soror* Forel and *Oecophylla smaragdina* Smith attended three aphid species each.

*Aphis craccivora* Koch and *A. spiraecola* Patch were attended by five species of ants, while *Rhopalosiphum maidis* (Fitch), *A. gossypii* and *Toxoptera odinae* (van der Goot) were attended by four species. *Hysteroneura setariae* (Thomas) and *T. aurantii* (Boyer de Fonscolombe) were attended by three species of ants and *T. citricida* (Kirkaldy), *Tetraneura nigriabdominalis* (Sasaki) and *Greenidea ficicola* Takahashi were attended by two ant species each. Of the 24 species of aphids attended by ants, majority of the species belonged to the genus *Aphis* (five), followed by *Toxoptera* (three) and *Greenidea* (two). *Oecophylla smaragdina* Smith attended aphids infesting citrus and mango, interestingly this species of ant is also considered as a pest on these host plants.

TABLE 1. List of aphid species recorded from Karnataka during the study

Subfamily	Tribe	Species
Aphidinae	Aphidini	<i>Aphis achyranthi</i> Theobald <sup>††</sup>
		<i>Aphis craccivora</i> Koch
		<i>Aphis fabae</i> Theobald
		<i>Aphis glycines</i> Matsumura <sup>†</sup>
		<i>Aphis gossypii</i> Glover
		<i>Aphis nerii</i> (Boyer de Fonscolombe)
		<i>Aphis punicae</i> Passerini
		<i>Aphis spiraecola</i> Patch
		<i>Hyalopterus pruni</i> (Geoffroy) <sup>††</sup>
		<i>Hysteroneura setariae</i> (Thomas)
		<i>Melanaphis donacis</i> (Passerini)
		<i>Melanaphis sacchari</i> (Zehntner)
		<i>Rhopalosiphum maidis</i> (Fitch)
		<i>Rhopalosiphum nymphaeae</i> (L.)
		<i>Rhopalosiphum padi</i> (Linn.) <sup>††</sup>
		<i>Rhopalosiphum rufiabdominalis</i> (Sasaki)
		<i>Schizaphis rotundiventris</i> (Signoret) <sup>††</sup>
		<i>Schizaphis graminum</i> (Rondani)
		<i>Toxoptera aurantii</i> (B. d. Fonsc.)
		<i>Toxoptera citricida</i> (Kirkaldy)
		<i>Toxoptera citricida</i> (Kirkaldy)
	Macrosiphini	<i>Acyrtosiphon gossypii</i> Mordvilko <sup>††</sup>
		<i>Acyrtosiphon kondoi</i> Shinji & Kondo
		<i>Acyrtosiphon pisum</i> (Harris)
		<i>Brachycaudus helichrysi</i> (Kalt.)
		<i>Brevicoryne brassicae</i> (Linn.)
		<i>Capitophorus elaeagni</i> (del Guercio)
		<i>Coloradoa rufomaculata</i> (Wilson)
		<i>Hayhurstia atriplicis</i> (Linnaeus) <sup>††</sup>
		<i>Hydaphis coriandri</i> (Das)
		<i>Hyperomyzus carduellinus</i> (Theobald) <sup>††</sup>
		<i>Ipuka dispersum</i> (van der Goot) <sup>††</sup>
		<i>Macrosiphoniella sanborni</i> (Gill.)
		<i>Macrosiphum rosae</i> (Linn.)
		<i>Myzus persicae</i> (Sulzer)
		<i>Pentalonia nigronervosa</i> Coquerel
		<i>Sitobion leelamantiae</i> David
		<i>Sitobion rosaeiformis</i> (Das)
		<i>Sitobion takahashii</i> Eastop
		<i>Uroleucon (Uromelan) compositae</i> (Theobald)
		<i>Uroleucon (Uroleucon) sonchi</i> (Linnaeus)

Contd...

TABLE 1. (Contd..)

Subfamily	Tribe	Species
<b>Chaitophorinae</b>		<i>Chaitophorus himalayensis</i> (Das) <sup>†</sup>
<b>Drepanosiphinae</b>		<i>Therioaphis trifolii</i> (Monell)
		<i>Unipterus</i> ( <i>Paoliella</i> ) <i>nirmalae</i> David
		<i>Tinocallis kahawaluokalani</i> (Kirkaldi) <sup>††</sup>
<b>Greenideinae</b>	Cervaphidini	<i>Cervaphis rappardi indica</i> Basu
		<i>Schoutedenia emblica</i> (Patel & Kulkarni)
	Greenideini	<i>Greenidea</i> ( <i>Greenidea</i> ) <i>artocarpi</i> (Westwood)
		<i>Greenidea</i> ( <i>Greenidea</i> ) <i>ficicola</i> Takahashi
		<i>Greenidea</i> ( <i>Trichosiphum</i> ) <i>annonae</i> (Pergande) <sup>††</sup>
		<i>Greenidea</i> ( <i>Trichosiphum</i> ) <i>formosana heeri</i>
		Raychaudhuri, Ghosh, Banerjee and Ghosh <sup>††</sup>
<b>Hormaphidinae</b>	Cerataphidini	<i>Astegopteryx bambucifoliae</i> (Takahashi)
		<i>Astegopteryx bambusae</i> (Buckton)
		<i>Astegopteryx formosana</i> (Takahashi)
		<i>Cerataphis brasiliensis</i> (Hempel) <sup>†</sup>
		<i>Cerataphis lataniae</i> (Boisduval)
		<i>Cerataphis orchidearum</i> (Westwood) <sup>††</sup>
		<i>Ceratovacuna lanigera</i> Zehntner
		<i>Pseudoregma bambusicola</i> (Takahashi)
<b>Lachninae</b>	Cinarini	<i>Cinara tujafilina</i> (Del Guercio) <sup>†</sup>
	Lachnini	<i>Tuberolachnus salignus</i> Gmelin <sup>†</sup>
<b>Pemphiginae</b>	Eriosomatini	<i>Forda orientalis</i> George
		<i>Tetraneura</i> ( <i>Tetraneurella</i> ) <i>nigriabdominalis</i> (Sasaki)
		<i>Tetraneura</i> ( <i>Tetraneurella</i> ) <i>javensis</i> Goot <sup>††</sup>

<sup>††</sup> First record from Karnataka<sup>†</sup> First record from south India

Earlier workers from south India indicated only presence or absence of ant attendance without mentioning the ant species involved with the exception of Gadiyappanavar (1970) who recorded *C. compressus* and *S. geminata* attending *T. nigriabdominalis*. Same species of ants were found to attend *T. nigriabdominalis* in the present study. In addition to this Musthak Ali and Sharatchandra (1986) and Veeresh and Musthak Ali (1990) have enlisted aphidocolous ants from southern India. More studies on aphid and ant relationship are available from northern and northeastern parts of India as compared to southern parts of India.

Twenty-four new aphid-ant associations (marked by an asterisk in second column of Table 2) were recorded in the present study. Studies thus indicated that the area of survey is rich in aphid as well as ant species attending to them.

TABLE 2. Aphidocolous ants (Hymenoptera: Formicidae) collected from Karnataka

Ant subfamily & species	Associated aphid species	Host plant
<b>Formicinae</b> <i>Camponotus</i> <i>comprassus</i> Fabricius	<i>Aphis craccivora</i> Koch	<i>Cajanus cajan</i> (Linn.) <i>Crotolaria juncea</i> Linn. <i>Crotolaria mucronata</i> Desvaux <i>Dolichos lablab</i> Linn. <i>Gliricidia maculata</i> (Jacq.) Kunth ex Walp. <i>Mimosa pudica</i> Mill. <i>Vigna unguiculata</i> (Linn.) Walp.
	<i>Aphis fabae</i> Theobald	<i>Cestrum nocturnum</i> Linn. <i>Solanum nigrum</i> Linn.
	<i>Aphis gossypii</i> Glover	<i>Solanum melongena</i> Linn. <i>Solanum</i> sp. <i>Gossypium hirsutum</i> Linn <i>Hibiscus rosa-sinensis</i> Linn.
	<i>Aphis spiraeicola</i> Patch *	<i>Chromolaena odorata</i> (Linn.) King & Robs
	<i>Cinara tujaefilina</i> (Del guercio)	<i>Thuja chilensis</i> Borders and Gaussen
	<i>Greenidea ficicola</i> Takahashi *	<i>Ficus tsiela</i> Roxb
	<i>Hysteroneura setariae</i> (Thomas)	<i>Cyperus rotundus</i> Miq. <i>Chlorophytum</i> sp. <i>Eleusine coracana</i> (Linn.) Gaertn.
	<i>Melanaphis sacchari</i> (Zenhnter)	<i>Sorghum bicolor</i> (Linn.) Moench.
	<i>Rhopalosiphum maidis</i> (Fitch)	<i>Zea mays</i> Linn.
	<i>Schoutedenia emblica</i> (Patel & Kulkarni) *	<i>Phyllanthus emblica</i> Linn.
	<i>Sitobion takahashii</i> Eastop *	<i>Phyllanthus emblica</i> Linn.
	<i>Tetraneura nigriabdominalis</i> (Sasaki)	<i>Eleusine coracana</i> (Linn.) Gaertn.
	<i>Toxoptera aurantii</i> (B.d.F.)	<i>Ixora</i> sp. <i>Camelia theifera</i> Griff
	<i>Toxoptera citricida</i> (Kirkaldy)	<i>Citrus</i> sp.
	<i>Toxoptera odinae</i> (van der Goot)	<i>Anacardium occidentale</i> Linn.
	<i>Ceratovacuna lanigera</i> Zehntner	<i>Saccharum officinarum</i> Salisb.
<i>Camponotus</i> <i>rufoglaucus</i> Jerd.	<i>Aphis craccivora</i> Koch	<i>Sesbania grandiflora</i> (Linn.) Pers
	<i>Aphis fabae</i> Theobald	<i>Cerstrum nocturnum</i> Linn. <i>Solanum nigrum</i> Linn.
<i>Crematogaster</i> <i>ransonneti</i> Mayr	<i>Tuberolachnus salignus</i> (Gmelin) *	<i>Salix</i> sp.
<i>Crematogaster</i> <i>soror</i> Forel.	<i>Aphis craccivora</i> Koch *	<i>Arachis hypogaea</i> Willd.
	<i>Aphis gossypii</i> Glover*	<i>Solanum melongena</i> Linn.
	<i>Aphis spiraeicola</i> Patch*	<i>Chromolaena odorata</i> (Linn.) King & Robs.

Contd...

TABLE 2. (Contd...)

Ant subfamily & species	Associated aphid species	Host plant
<i>Oecophylla smaragdina</i> Smith	<i>Chaitophorus himalayensis</i> (Das)*	<i>Salix</i> sp.
	<i>Toxoptera aurantii</i> (B.d.F.)	<i>Ixora</i> sp. <i>Citrus</i> sp.
	<i>Toxoptera odinae</i> (van der Goot)*	<i>Hibiscus rosa-sinensis</i> Linn. <i>Mangifera indica</i> Linn.
<b>Myrmicinae</b>		
<i>Monomorium gracillium</i> Smith	<i>Unipterus nirmalae</i> David *	<i>Terminalia arjuna</i> (Roxb.ex DC.) Wight & Arn.
<i>Monomorium</i> sp.	<i>Aphis affinis</i> Del Guercio *	<i>Mentha viridis</i> (Linn.)
	<i>Aphis craccivora</i> Koch	<i>Vigna unguiculata</i> (Linn.) Walp.
	<i>Aphis gossypii</i> Glover	<i>Gossypium hirsutum</i> Linn.
	<i>Aphis spiraeicola</i> Patch	<i>Chromolaena odorata</i> (Linn.) King & Robs.
	<i>Rhopalosiphum maidis</i> (Fitch)	<i>Zea mays</i> Linn.
	<i>Toxoptera aurantii</i> (B.d.F.) *	<i>Camelia theifera</i> Griff
	<i>Toxoptera odinae</i> (van der Goot) *	<i>Anacardium occidentale</i> Linn. <i>Hibiscus rosa-sinensis</i> Linn. <i>Ixora</i> sp.
	<i>Greenidea</i> ( <i>Trichosiphum</i> ) <i>formosana heeri</i> Raychaudhuri <i>et al.</i> *	<i>Psidium guajava</i> Linn.
	<i>Dactynotus sonchi</i> Linnaeus	<i>Sonchus arvensis</i> Linn.
<i>Myrmicaria brunnea</i> Saunders	<i>Rhopalosiphum maidis</i> (Fitch) *	<i>Zea mays</i> Linn.
<i>Solenopsis geminata</i> Fabricius	<i>Aphis craccivora</i> Koch	<i>Vigna unguiculata</i> (Linn.) Walp. <i>Dolichos lablab</i> Linn. <i>Gliricidia maculata</i> (Jacq.) Kunth ex Walp.
	<i>Aphis nerii</i> (B.d.F.) *	<i>Asclepias curassavica</i> Griseb. <i>Asclepias physocarpa</i> Schlechter
	<i>Aphis spiraeicola</i> Patch *	<i>Chromolaena odorata</i> (Linn.) King & Robs.
	<i>Pentalonia nigrinervosa</i> Coquerel	<i>Musa sapientum</i> Linn. <i>Alpinia versicolor</i> K.Schum. <i>Maranta zebrina</i> Sims
	<i>Rhopalosiphum maidis</i> (Fitch) *	<i>Zea mays</i> Linn.
	<i>Hysteroneura setariae</i> (Thomas) *	<i>Eleusine coracana</i> (Linn.) Gaertn.
	<i>Tetraneura niglabdominalis</i> (Sasaki)	<i>Eleusine coracana</i> (Linn.) Gaertn.
	<i>Toxoptera citricida</i> (Kirkaldy) *	<i>Citrus</i> sp.
<b>Dolichoderinae</b>		
<i>Tapinoma melanocephalum</i> Fabricius	<i>Aphis spiraeicola</i> Patch *	<i>Chromolaena odorata</i> (Linn.) King & Robs.
	<i>Hysteroneura setariae</i> (Thomas) *	<i>Eleusine coracana</i> (Linn.) Gaertn.
<i>Technomyrmex albipes</i> Smith	<i>Aphis gossypii</i> Glover *	<i>Hibiscus rosa-sinensis</i> Linn.
	<i>Toxoptera odinae</i> (van de goot) *	<i>Hibiscus rosa-sinensis</i> Linn.

\*New aphid ant associations

## ACKNOWLEDGEMENTS

I am indebted to Dr. C. A. Viraktamath, Scientist Emeritus, University of Agricultural Sciences, GKVK, Bangalore for guidance, technical support and comments on the paper. I am thankful to Prof. T. M. Musthak Ali, Professor, Department of Entomology, UAS, GKVK, Bangalore for identification of ant species. The study leave granted to me by the Project Director, Project Directorate of Biological Control (ICAR), Hebbal, Bangalore, is gratefully acknowledged.

## REFERENCES

- Addicott, J.F. (1978) Competition for mutualists: aphids and ants. *Can. J. Zool.* **56**: 2093–2096.
- Banks, C.J. (1958) Effects of ant, *Lasius niger* (Linn.) on the behaviour and reproduction of the black bean aphid, *Aphis fabae* Scopoli. *Bulletin of Entomological Research* **49**: 701–714.
- Banks, C.J. and Macaulay, E.D.M. (1967) Effects of *Aphis fabae* Scopoli and its attendant ants and insect predators on yields of field beans (*Vicia faba* L.). *Ann. Appl. Biol.* **60**: 445–453.
- Banks, C.J. and Nixon, H.L. (1958) Effects of the ant *Lasius niger* (L.) on the feeding and excretion of the bean aphid, *Aphis fabae* Scopoli. *J. Exp. Biol.* **35**: 703–711.
- Bisht, R.S., Rana, D.S. and Katoch, A.R. (2002) Records of aphidicolous ants (Hymenoptera: Formicidae) from Garhwal hills, Utranchal. *J. Aphidol.* **16**: 15–19.
- Broadbent, L. (1951) Aphid excretion. *Proc. Res. Ent. Soc. London (A)* **26**: 97–103.
- Curtright, C.R. (1965) Subterranean aphids of Ohio. *Bull. Ohioagric. Expt. Stn* **387**: 175–233.
- Datta, S.K., Raychaudhuri, D. and Agarwala, B.K. (1982) Study on aphid tending ants in India-I. *Entomon* **7**: 327–328.
- Datta, S.K., Raychaudhuri, D. and Agarwala, B.K. (1983) Study on aphid tending ants in India-II. *Entomon* **8**: 23–25.
- Datta, S.K. and Raychaudhuri, D. (1984) New aphidicolous ants (Hymenoptera: Formicidae) from India. *Newsl. Aphidol. Soc. India* **4**: 7.
- Datta, S.K. and Raychaudhuri, D. (1985) Taxonomy of aphidicolous ants (Hymenoptera: Formicidae) of Sikkim and hilly areas of West Bengal. *Akitu* **56**: 1–14.
- Devi, C.M. and Singh, T.K. (1986a.) New aphidicolous ants from India. *Newsl. Aphidol. Soc. India* **5**: 5–7.
- Devi, C.M. and Singh, T.K. (1986b.) Aphid tending ants on brinjal plant (*Solanum melongena*) in Manipur. *Newsl. Aphidol. Soc. India* **5**: 7.
- Devi, C.M., Paonam, M.S. and Singh, T.K. (1987) Aphids (Homoptera: Aphididae) and their associated ants (Hymenoptera: Formicidae) of fruit trees in Manipur. *J. Aphidol* **1**: 102–104.
- Devi, C.M., Singh, T.K. and Nilamani, L. (2001) New aphidicolous ants (Hymenoptera: Formicidae) from India: Subfamily: Myrmicinae. *J. Aphidol* **15**: 107–112.
- Eastop, V.F. (1966) A taxonomic study of Australian Aphidoidea. *Australian Journal of Zoology* **14**: 399–592.
- El-Ziady, S. (1960) Further effects of *Lasius niger* Linn. on *Aphis fabae* Scopoli. *Proc. R. Ent. Soc. London. (A)* **35**: 30–38.
- Flanders, S.E. (1951) The role of the ant in the biological control of homopterous insects. *Canadian Entomologist* **83**: 93–98.
- Gadiyappanavar, R.D. (1970) Study of aphid pests of crops of Mysore State and the biology of ragi root aphid, *Tetraneura nigriabdominalis* (Homoptera: Aphididae) *M.Sc. Thesis*, Submitted to the University of agricultural Sciences, Bangalore.
- Gadiyappanavar, R.D. (1970) Faunistic study of aphids of Mysore and the biology of ragi root aphid, *Tetraneura nigriabdominalis* (Sasaki) (Homoptera: Aphididae) *M.Sc. Thesis*.



- Submitted to Department of Agricultural entomology, University of Agricultural Sciences, Bangalore, p. 110, (unpublished)
- Ghosh, A.K. (1986) Aphidological studies in India. In: *Recent Trends in aphidological Studies*, Kurl. S.P. (Ed). Aphidological Society of India, Modinagar, 1–6.
- Krishnamurti, B. (1928) Aphididae of Mysore, I. *Journal of Bombay Natural History Society* **33**: 211–215.
- Krishnamurti, B. (1930) Aphididae of Mysore, II. *Journal of Bombay Natural History Society* **34**: 411–419.
- Krishnamurti, B. (1950) Aphididae of Mysore, III. *Indian Journal of Entomology* **10**: 51–53.
- Kurl, S.P. and Misra, S.D. (1980) Interaction between the aphid and their attended ants. *J. Env. Res.* **1**: 1–6.
- Martin, J.H. (1983) The identification of common aphid pests of tropical agriculture. *Tropical Pest Management* **29**: 395–411.
- Matta, T.A. and Bhagat, R.C. (2000) Diversity of aphidocolous ants in Kashmir with reference to ant-aphid association. In: *Environment, Biodiversity and Conservation*, Khan, M.A. (Ed). A.P.H. Publishing Corporation, 532.
- Musthak Ali, T.M. and Sharatchandra, H.C. (1986) Role of ants in establishment of root aphid *Forda orientalis* George on finger millet *Eleusine coracana* Beauv and its management. *Proceedings of the National Symposium on Aphidology in India II held at Agaratala 2-4 Nov. 1985* 75–80.
- Nieto Nafria, J.M., Mier Durante, M.P. and Remaudière, G. (1998) Catalogo de los Pulgones de España y de sus Plantas Hospedadores. Universidad de León (Secretariado de Publicaciones), León (Spain), 17p
- Raychaudhuri, D.N. (1980) *Aphids of North-East India and Bhutan*, Zoological Society, Calcutta, 521p.
- Remaudière, G. and Remaudière, M. (1977) *Catalogue des Aphididae du Monde*, INRA, Paris, 473.
- Roy, D.K. and Behura, B.K. (1980) Three more species of ants attending *Aphis gossypii* Glover. *Newsl. Aphidol. Soc. India* **1**: 4.
- Shantibala., Singh, T.K. and Varathanjan, R. (1995) Studies on the association between *Cervaphis quercus* Takahashi (Homoptera: Aphididae) and its tending ants. *J. Aphidol.* **9**: 55–62.
- Skinner, G.J. and Whittaker, J.E. (1981) An experimental investigation of inter-relationships between the wood ant (*Formica rufa*) and some tree-canopy herbivores. *J. Anim. Ecol.* **50**: 313–326.
- van Emden, H.F. (1972) *Aphid Technology*, Academic Press, London. xiv + 344.
- Veeresh, G.K. and Musthak Ali, T.M. (1990) The present status of our knowledge on Indian ants. In: *Social Insects Perspective*, Veeresh, G.K., Kumar, A.R.V. and Shivashankar, T. (Eds). IUSSI-Indian Chapter, Bangalore, India.
- Verghese, A. and Tandon, P.L. (1987) Interspecific associations among *Aphis gossypii*, *Menochilus sexmaculatus* and *Camponotus compressus* in a guava ecosystem. *Phytoparasitica* **5**: 289–297.
- Way, M.J. (1963) Mutualism between ants and honeydew producing Homoptera. *Ann. Rev. Ent.* **8**: 307–344.

(Received 21 December 2007; accepted 15 February 2008)





## Voltage gated potassium channels in the larval CNS neurons of coconut black headed caterpillar, *Opisina arenosella* Wlk.

K. Subaharan<sup>\*1</sup> and Murali Mohan<sup>2</sup>

<sup>1</sup>Central Plantation Crops Research Institute, Kasaragod 671124, Kerala, India  
Email: subaharan\_70@yahoo.com

<sup>2</sup>Dept. of Entomology, UAS, GKVK Campus, Bangalore, Karnataka, India

**ABSTRACT:** A primary dissociated cell culture of *Opisina arenosella* Wlk. larval Central Nervous System (CNS) neurons was established. The acutely dissociated larval CNS neurons could be divided into two categories based on their morphology and size. Type I consisted of rounded cells and type II consisted of cells with processes. Nearly 60 per cent of the cells were alive after 7 days of plating. Whole cell currents were recorded from Type II cells (5–12  $\mu\text{m}$ ) of 12–48 h using the patch clamp setup. The outward potassium currents were obtained from the neuronal cells. The absence of potassium currents from the cells when exposed to potassium channel blockers (cocktail of 4-aminopyridine (1 mM), quinidine (0.1 mM) and tetraethylammonium (TEA, 20 mM) in extra cellular bath) confirmed the potassium currents. Three types of potassium currents (non inactivating, slow inactivating, and fast inactivating currents) were observed in whole cell recording of Type II cells.

© 2008 Association for Advancement of Entomology

**KEYWORDS:** *Opisina arenosella*, cell culture, patch clamp, potassium currents

### INTRODUCTION

Neuronal signaling depends on rapid change in the electrical potential differences across nerve cell membranes. Insect neuronal cell membranes have a variety of distinct potassium channels that can be activated by changes in membrane voltage (Latorre *et al.*, 1984). The relative number and spatial distribution of these channels in the cell determine the particular excitability properties of the neuron (Adams *et al.*, 1980; Thompson and Aldrich, 1980). The information about voltage-dependant ion channels (VDIC) have been investigated using cultured neurons of *Drosophila melanogaster* (Wu and Suzuki, 1983; Subaharan, 2004) and other insects (Hayashi and Hildebrand, 1990). Ion channels are primary target site for several classes of

\*Corresponding author

natural and synthetic insecticide compounds (Okuma and Yoshiyuki, 2004). Potent natural toxins like apamin, and dendrotoxin and ecdysteroid agonist RH-5849 block the potassium channels (Castle *et al.*, 1989; Salgado, 1992a,b).

The coconut black headed caterpillar, *Opisina arenosella* Wlk. is a serious pest on coconut (Singh and Rethinam, 2006). The larvae feed on the parenchymatous tissues on the undersurface of the coconut leaflets and constructs galleries of silk and excreta (Nirula *et al.*, 1951). This pest occurs in Kerala, Karnataka, Tamil Nadu, Andhra Pradesh, Orissa, West Bengal, Maharashtra and Gujarat states of India (Venkatesan *et al.*, 2003). Though the inundative release of parasitoids keeps the population of the pest under check their timely availability for release during the outbreak is a limiting factor. Hence, during the outbreak it warrants the use of pesticides to check the spread of pest to newer locations. Currently farmers resort to spraying of malathion, dichlorvos (Ponnama, 1984) and stem injection or root feeding of monocrotophos (Shivakumar, 1999; Patalappa, 1988). As monocrotophos and dichlorvos have been brought under the restricted use pesticide status, there is a need to search for insecticides with novel chemistry that can be harmoniously used in integrated management of this pest. During the process of screening of newer molecules bioassay coupled with electrophysiology studies would furnish additional information at cellular level on the potency of neurotoxic insecticides. This study can be facilitated by establishing dissociated neuronal cultures of the test insect and then studying its channel properties. Examining the K<sup>+</sup> channels in *O. arenosella* neuronal membrane by biophysical methods would provide a powerful approach for studying the channel properties.

No earlier attempts have been made to culture CNS neurons from *O. arenosella* larvae. As *in vitro* cell culture allows continuous observation on isolated cells, an attempt was made to establish a primary culture of dissociated neurons of CNS of seventh instar *O. arenosella* larval brain. The suitability of dissociated CNS cell culture of *Drosophila melanogaster* for patch clamp analysis has been suggested (Sun and Wu, 1984, Solc and Aldrich, 1988, Yamamoto and Suzuki, 1987). Hence, in the present study we attempted to develop a dissociated CNS neuronal cell culture of *O. arenosella* and characterize the voltage dependent K channels at the whole cell level by patch clamp technique.

## MATERIALS AND METHODS

### Insect culture

The larvae of *O. arenosella* were collected from the Tiptur tall palms in Tumkur block of Karnataka State in India. The insects were reared in the laboratory in a glass jar provided with fresh coconut leaves, following the method of Santhosh Babu and Prabhu (1987).

### Cell culture

To obtain the CNS neurons, uniform size seventh instar *O. arenosella* larvae were taken and rinsed in double distilled water twice followed by a rinse in 70 % ethanol for 1–2 min. They were again rinsed in double distilled water to remove the traces of alcohol. The larval brain was dissected using micro forceps in phosphate buffer saline (PBS) and was immediately transferred to another dish containing PBS to prevent the contamination due to regurgitated material by the larvae while dissecting. The brain was then mechanically teased in PBS and then incubated in  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  free PBS containing 0.5 mg/ml collagenase (Type I Sigma) saline for one hour in an eppendorf tube in sterile condition. After incubation the contents in the tube were triturated 50 times with a pipette tip. The preparation was centrifuged for 1–2 min at 3000 rpm. The supernatant was discarded. The pellet was re-suspended in Grace insect medium (Sigma) supplemented with 1.5% penicillin/streptomycin (Sigma) and 10 % Fetal Bovine Serum (Gibco). The cell suspension was plated on autoclaved cover slips at a density of one brain complex / cover slip. The cover slips were coated with Poly L Lysine (Sigma). The cover slips were then placed in a sterile 35 mm Petri dish (Tarsons) and incubated at 19–22 °C in a humidified incubator. The cell types were also studied based on the morphology.

In order to assess the viability of CNS neurons, the cell culture was observed at varied intervals (1, 2, 4, 6, 8 and 10 days after plating) under an inverted phase contrast microscope (Nikon–Diaphot 200). Each time destructive sampling was done using Tyrosine blue stain as indicator. The per cent mortality of cells at varied intervals was worked out.

### Electrophysiology

Membrane currents of neurons were studied in order to characterize the type of channels present in the CNS neurons. Recordings were made from 2–6 days old dissociated CNS neurons cell culture having 5–12  $\mu\text{m}$  long processes. Currents were recorded using the whole-cell configuration of the patch-clamp technique using an Axopatch amplifier (Axon instruments). The patch-clamp electrodes with a resistance of 2–8  $\text{M}\Omega$  were fabricated from borosilicate glass with a Narishige PP-83 puller. The capillaries were polished with Narishigae microforge. The patch pipettes of around 1  $\mu\text{m}$  diameter were filled with internal solution.

Whole cell currents were recorded from 2, 4 and 6 day old bipolar neurons (5–12  $\mu\text{m}$ ), using the patch clamp setup (Axon instruments). In external bath solution, a typical electrode resistance of around 2–8  $\text{M}\Omega$  was obtained. The seal was made by a brief suction at the cell surface after the electrode touched the cell. A tight seal was immediately formed and a typical resistance of around 1–50  $\text{G}\Omega$  was obtained. The capacitive transients were minimized. A whole cell configuration was attained by a few light suctions. Increase in the size of capacitive transients and a little fall in the seal resistance confirmed whole cell configuration. Capacitive transients were compensated at this stage. The cells were voltage clamped at  $-100\text{ mV}$  and

an increasing step of 20mV was given each time (i.e.  $-80$  mV,  $-60$  mV,  $-40$  mV,  $-20$  mV,  $0$  mV,  $20$  mV,  $40$  mV,  $60$  mV, and  $80$  mV) for 200 ms interval.

The recordings were acquired and analyzed using *pClamp6.0* software (Axon Instruments) run on an IBM compatible '386' computer. Currents recorded from cells were filtered online using 1000 Hz frequency filter. After acquisition, the baseline of the currents was compensated and the leak subtracted. Amplitude of the peak currents obtained in the cultures was measured.

## RESULTS

### Dissociated neurons of *Opisina arenosella* CNS in primary cell cultures

After plating, the dissociated CNS neuronal cells settled down and got attached to the poly-lysine coated dishes within 3 h of plating. The cells, which do not adhere to the dish, remain rounded and died out. There were no glial cells in the culture.

The acutely dissociated larval CNS neurons could be divided into two categories based on their morphology and size. Type I consisted of rounded cells (diameter  $> 5 \mu\text{m}$ ) with no process and they comprised approximately 30% of the population. Type II cells comprised approximately 70% of the total population. In addition to soma they had ( $5\text{--}7 \mu\text{m}$ ) processes (Fig. 1).

### Viability of CNS neuronal cells in culture

Viability of the cells was measured as a function of days using hemocytometer. Trypan blue permeable cells were counted as dead. More than 60% of the cells were alive 6 days after plating (Table 1).

### Heterogeneity of whole cell currents

The acutely dissociated CNS neuronal type I cells were not suitable for forming a giga ohm seal during whole cell recording. Hence, the type II cells, which comprised the largest population of dissociated CNS neuronal cells, were selected for the study of membrane currents.

The currents obtained in all the larval CNS neurons clamped exhibited only net outward currents in response to depolarizing step pulses ( $-80$ ,  $-60$ ,  $-40$ ,  $-20$ ,  $0$ ,  $20$ ,  $40$ ,  $60$ ,  $80$  mV). The outward currents were similar to potassium currents. A cocktail of 4-aminopyridine (1 mM), Quinidine (0.1 mM) and tetraethylammonium (TEA, 20 mM) was used in extra cellular bath to block all the currents passing through potassium channels as suggested by Deepti (2002). No currents were observed in the presence of these blockers. This confirmed that the outward currents were potassium channel specific. Also, in absence of potassium currents no other type of currents could be observed in cells in culture. The current/voltage relationship was measured for the various depolarizing voltage commands.

Heterogeneity of the whole cell outward currents in peak amplitude and inactivation time constants existed. The classification of  $\text{K}^+$  currents was the first step towards



FIGURE 1. Dissociated *O. arenosella* larval CNS neuronal cell culture. Black arrow shows the Type I spherical cells and white open arrow shows the Type II cells with processes.

TABLE 1. Viability of neuronal cells

Age of the neuronal cells (days)	Per cent mortality $\pm$ SE
1	$92.2 \pm 1.06^a$
2	$90.2 \pm 0.58^{ab}$
4	$86.8 \pm 1.15^b$
6	$63.6 \pm 2.73^c$
8	$53.4 \pm 2.80^c$
10	$37.6 \pm 2.50^c$

Means with same letter are not significantly different at  $p = 0.05$  with DnMRT.

detecting an alteration in  $K^+$  currents. Three kinds of outward  $K^+$  currents were observed. This classification was according to their fast inactivation time constants. All the three types of currents were voltage dependent and their I/V curves were similar (Fig. 2—4). Some cells exhibited primarily a fast transient K current that turned on and inactivated rapidly in response to depolarizing voltage steps whilst some neurons had K currents with slower inactivation time course. A group of cells also expressed a non inactivating component of outward current.

### Types of potassium currents

Three kinds of voltage sensitive potassium currents were distinguishable. These were classified on the basis of their activation and inactivation kinetics.

#### *Non inactivating type*

No inactivation of the channels were observed in 200 ms. Peak current was obtained after  $80.80 \pm 1.40$  ms of the beginning of the voltage step. The activation of K1




FIGURE 2. Non inactivating type currents: The traces here show the  $-80$ ,  $-60$ ,  $-40$ ,  $-20$ ,  $0$ ,  $20$ ,  $40$ ,  $60$ ,  $80$  mv pulses after the baseline correction and leak subtraction. The plot of current vs voltage is on the right.




FIGURE 3. Slow inactivating currents. The traces here show the  $-80$ ,  $-60$ ,  $-40$ ,  $-20$ ,  $0$ ,  $20$ ,  $40$ ,  $60$ ,  $80$  mv pulses after the baseline correction and leak subtraction. The plot of current vs voltage is on the right.






FIGURE 4. A fast inactivating current. The traces here show the  $-80$ ,  $-60$ ,  $-40$ ,  $-20$ ,  $0$ ,  $20$ ,  $40$ ,  $60$ ,  $80$  mv pulses after baseline correction and leak subtraction. The plot of current vs voltage is on the right.

TABLE 2. Types of K<sup>+</sup> currents in neuronal culture

Days after plating	Number of cells showing (currents)		
	Fast inactivating	Slow inactivating	Non inactivating
2	60.00	10.00	30.00
4	58.13	11.62	30.23
6	55.55	16.66	27.77

channels followed first order exponential kinetics with time constant of activation  $20.48 \pm 1.6$  ms (Fig. 2).

#### *Slow inactivating type*

These channels activate faster than K1 channels with peak current at  $26.80 \pm 2.0$  ms. The inactivation kinetics is slow and follows a single exponential function of inactivation (Fig. 3).

#### *Fast inactivating type*

These are fast inactivation type channels. The activation of these channels after the voltage pulse beginning was very fast and the peak current reached at around 12.2 to 4.77 ms. The inactivation was also very fast. Two time constant might reflect the intrinsic property of these channels (Fig. 4).

Analysis of the decay phase of the currents recorded from larval CNS neurons of *O. arenosella* showed the following distribution. Irrespective of the age majority of the cells showed the fast transient currents followed by non inactivating or sustained currents. A minority of the neurons exhibited primarily a delayed transient K<sup>+</sup> currents (Table 2). There was not much difference in the type of potassium currents with increase in age of the cells.

## DISCUSSION

The primary culture of dissociated CNS neurons from VII instar *O. arenosella* larvae was established. Two types of neurons were observed in cultures. Majority of the cells had processes. In case of *D. melanogaster*, three types of cells were observed in the dissociated larval CNS culture. They were differentiated based on their size and morphology (Wu *et al.*, 1983; Subaharan, 2004). Sang (1981) reported that neuronal cultures obtained from larval brains by dissection was relatively free from glial cells. In our studies also we observed that the frequency of cells with non neuronal morphology was less than 1% in cultures of different ages. The relatively low number of glial cells in culture was not due to its inability to survive in culture medium but due to the lower number of glial cells in insect CNS.

In the dissociated CNS cell culture more than 60 per cent of the cells were viable 4 days after plating. Neuronal cultures have proved to be useful for a number of studies. The neuronal cells in the culture system aids in the study of membrane currents by electrophysiological technique (Wu *et al.*, 1983). Primary cultures have been used to record neuronal current properties owing to the easily manageable conditions in cultures as compared to *in vivo*. Primary cultures have been previously made from central nervous system of *Drosophila* (Wu *et al.*, 1983).

There have been several studies on voltage dependant current in insect neurons (Leung and Byerly, 1991; Saito and Wu, 1991; Solc and Aldrich, 1988). No studies have been done on *O. arenosella*. Hence our goal was to examine the membrane currents of the cultured CNS neuronal cells. Patch clamp recordings were done on neurons from VII instar *O. arenosella* larval CNS cultures.

Potassium currents in larval CNS neurons had three components: a fast transient component, a more slowly inactivating component and a non-inactivating component. The relative amount of each component varied from cell to cell. Solc and Aldrich (1988) reported that the *D. melanogaster* larval CNS neurons had three distinct K channels; in addition they did not observe any voltage dependent Na currents in the cells studied. Diversity of K channel provides a basis for a wide spectrum of physiological properties among the cells recorded. K currents with diverse characters play a vital role in several phenomena such as depolarization of membrane potential, repetitive firing and sensory receptor potential (Hille, 1992).

Among the types of potassium currents recorded by whole cell recording majority of the channels; exhibited fast activating type i.e A type channel. The information obtained on the potassium channel on the neuronal membranes of *O. arenosella* larvae can be used to screen novel compounds that could be effective in their

management. Insecticides (Beleaf, FMC) with novel chemistry targets on the A type potassium channel (fast activating type) of central nervous system of the insect. These insecticides cause rapid and irreversible feeding cessation in the targeted pests, ultimately resulting in starvation and death.

#### ACKNOWLEDGEMENTS

The first author acknowledges Prof. Obaid Siddiqi for providing the facility of electrophysiology units for the study.

#### REFERENCES

- Adams, D. J., Smith, S. J. and Thompson, S. H. (1980) Ionic currents in molluscan soma. *Annual Review of Neuroscience* **3**: 141–167.
- Castle, N. A., Haylett, D. G. and Jenkinson, D. H. (1989) Toxins in the characterization of potassium channels. *Trends in Neuroscience* **12**: 59–65.
- Deepti, Trivedi (2002) Characterization of cshB : A chemosensory gene of *Drosophila melanogaster* M.Sc Thesis, MAHE, Manipal.
- Okuma, Eiji and Murata, Yoshiyuki (2004) Plant ion channels as potential targets for agro chemicals. *Journal of Pesticide Science* **29**(4): 304–307.
- Hayashi, J. H. and Hildebrand, J. G. (1990) Insect central olfactory neurons in primary culture. *Journal of Neuroscience* **10**: 848–859.
- Hille, B. (1992) Ionic channels of excitable membranes. Sinauer, Sunderland.
- Latorre, R., Coronado, R. and Vergara, C. (1984) K<sup>+</sup> channels gated by voltage and ions. *Annual Review of Physiology* **46**: 485–495.
- Leung, H. T. and Byerly, (1991) Characterization of single calcium channels in *Drosophila* embryonic nerve and muscle cells. *Journal of Neuroscience* **11**: 3047–3059.
- Patalappa, (1988) The control of coconut black headed caterpillar, *Opisina arenosella* by rot feeding. *Lal baugh* **30**(3): 3–6.
- Ponnama, K. N. (1984) Field studies on the comparative efficacy of BHC, dichlorvos and trichlorophan for the control of *Opisina arenosella* Wlk. *Indian coconut Journal* **15**(7): 3–5.
- Nirula, K. K., Antony, J. and Menon, K. P. V. (1951) Investigations on the pests of coconut palm, the coconut caterpillar, *Nephantis serinopa* Meyr. Life history and habits. *Indian Coconut Journal* **4**(4): 217–223.
- Saito, M. and Wu, C. F. (1991) Expression of ion channels and mutational effects in Giant *Drosophila* neurons differentiated from cell division arrested embryonic neuroblasts. *Journal of Neuroscience* **11**: 2135–2150.
- Salgado, V. L. (1992a) The neurotoxic insecticidal mechanism of non steroidal ecdysone agonist RH–5849: K<sup>+</sup> channel block in nerve and muscle. *Pesticide Biochemistry and Physiology* **43**: 1–13.
- Salgado, V. L. (1992b) Block of voltage dependant K channels in insect muscle by the diacylhydrazie insecticide RH–5849, 4-aminopyridine and quinidine. *Archives of Insect Biochemistry and Physiology* **21**: 239–252.
- Sang, J. H. (1981) *Drosophila* cells and cell lines. *Advances in Cell Culture* **1**: 125–183.
- Santhosh Babu, P. S. and Prabhu, V. K. K. (1987) How many larval instars are there in *Opisina arenosella*. *Entomon* **12**: 39–42.
- Solc, K. C. and Aldrich, W. R. (1988) Macroscopic and single-channel potassium currents in dissociated *Drosophila* CNS neurons. *Soc. Neurosci. Abstr.* **12**: 44.
- Singh, S. P. and Rethinam, P. (2006) Introduction to the coconut leaf eatingcaterpillar. *Opisina arenosella* Walker. *CORD* **22**: 1–24.

- Sun, Y. A. and Wu, C. F. (1984) Voltage dependant single-channel currents in dissociated CNS neurons of *Drosophila*. *Soc. Neurosci. Abstr.* **10**: 1091.
- Subaharan, K. (2004) Olfaction in *Drosophila*. Report submitted to DBT, Govt. of India.
- Shivakumar, T. (1999) New syringe method of insecticide administration technique to control coconut black headed caterpillar, *Opisina arenosella* Walker. *Mysore Journal of Agriculture Sciences* **33**(4): 301–305.
- Thompson, S. H. and Aldrich, R. W. (1980) Membrane potassium channels. In: *In the Cell Surface and Neuronal Function*, Cotman, C. W., Poste, G. and Nicolson, G. L. (Eds). Elsevier, North Holland Biomedical, Amsterdam, 49–85.
- Venkatesan, T., Jalali, S. K., Murthy, K. S., Rabindra, R. J. and Rao, N. S. (2003) A novel method of field release of *Goniozus nephantidis* an important parasitoid of *Opisina arenosella* on coconut. *Journal of Biological Control* **91**(1): 79–80.
- Woodward, C., Huang, T., Sun, H., Helfand, S. L and Carlson, J. R. (1989) Genetic analysis of olfactory behavior in *Drosophila melanogaster*: A new screen yields the *ota* mutants. *Genetics* **123**: 315–326.
- Wright, J. D. N. and Yi Zhong, (1995) Characterization of K currents and the CAMP-dependent modulation in cultured *Drosophila* mushroom body neurons identified by lac Z expression. *Journal of Neuro Science* **15**(2): 1025–1034.
- Wu, C. F., Suzuki, N. and Poo, M. M. (1983) Dissociated neurons from normal and mutant *Drosophila* larval central nervous system in cell culture. *Journal of Neuro science* **3**: 1888–1899.
- Wu, C. F. and Suzuki, (1983) Dissociated neurons from the normal and mutant *Drosophila* central nervous system in cell culture. *Journal of Neuro science* **3**: 1888–1899.
- Yamamoto, D. and Suzuki, N. (1987) Blockage of chloride channels by HEPES buffer. *Proceedings of Royal Society of London (Biol.)* **230**: 93–100.

(Received 5 December 2007; accepted 15 February 2008)



## Fecundity and developmental stages of Rice Blue Beetle, *Leptispa pygmaea* Baly (Coleoptera: Chrysomelidae) on rice varieties

K. Karthikeyan<sup>1</sup> and Sosamma Jacob<sup>2</sup>

<sup>1</sup>Regional Agricultural Research Station, Pattambi 679306, Kerala, India

Email: profkarthi@yahoo.co.in

<sup>2</sup>College of Horticulture, Agricultural University, Vellanikara 680656, Kerala, India

**ABSTRACT:** Biological studies on fecundity and developmental stages of rice blue beetle, *Leptispa pygmaea* Baly (Coleoptera :Chrysomelidae) were conducted at the Regional Agricultural Research Station Pattambi, Kerala Agricultural University, under net house condition on two rice varieties viz., Jyothi (short duration) and Aiswarya (medium duration) during June to October 2005. Adult beetle laid oval shaped yellow eggs on both the leaf surfaces either singly or in parallel rows with an average fecundity of 16.8 eggs on short duration variety Jyothi and 14.3 eggs on medium duration rice variety Aiswarya. The incubation period was 3.4 days in both varieties. The grub had five larval instars each with duration of 1–2 days and completed grub development within a mean period of 8.2 days. The grub pupated on the surface of leaf by getting attached loosely at its posterior end. The pupal period was 3.2 and 2.9 days in Jyothi and Aiswarya, respectively and completed the life cycle within 14.8 in Jyothi 13.8 days in Aiswarya. The pre-oviposition period of *L. pygmaea* showed no marked difference between varieties of rice. The longevity of adult varied with the sex of blue beetle. Males lived longer than females. Adult male of *L. pygmaea* lived for 40.9 days on Jyothi and 36.7 days on Aiswarya. The male life span was longer in Jyothi than Aiswarya. The females lived for an average period of 25 days in both varieties, thus indicating a shorter life for females than males.

© 2008 Association for Advancement of Entomology

**KEYWORDS:** *Leptispa pygmaea*, rice varieties, development, fecundity

### INTRODUCTION

The scenario of insect pest complex in rice has undergone a tremendous change during the past years. Certain insect pests of relatively minor importance have started appearing as major problems in rice. One among them is Rice Blue Beetle, *Leptispa pygmaea* Baly (Coleoptera: Chrysomelidae) which is becoming a serious problem in several rice growing districts of Kerala. The present investigations were, therefore,

undertaken to study the biological events of the pest under Kerala conditions. The study of life cycle of a pest is essential in order to identify the susceptible life stages which provide important clues for intervention and management.

#### MATERIALS AND METHODS

The biological studies of *Leptispa pygmaea* were carried out in the net house at the prevailing conditions of maximum temperature ( $30.1^{\circ}\text{C} \pm 1.40^{\circ}\text{C}$ ), minimum temperature ( $23.1^{\circ}\text{C} \pm 0.69^{\circ}\text{C}$ ) and relative humidity ( $94.33\% \pm 2.11\%$ ) during June to October, 2005. The biology was studied on two rice varieties viz., Jyothi (short duration) and Aiswarya (medium duration) by raising them in mud pots in the net house at the Regional Agricultural Research Station, Pattambi, Kerala Agricultural University. Adult beetles of *L. pygmaea* were collected from unsprayed rice fields and reared them on potted rice seedlings (variety Jyothi) covered with polyester cages of size  $49 \times 18$  cm in the net house. This served as a stock culture for the steady supply of beetles required for conducting various net house studies.

A pair of freshly mated beetles of *L. pygmaea* collected from the stock culture was released on 15 days old potted rice seedlings and covered with a polyester cage of size  $49 \times 18$  cm closed by muslin cloth at one end which served as a replication. Ten such replications were maintained on Jyothi and Aiswarya for studying the life cycle of *L. pygmaea*. Observations on the site of oviposition, pre-oviposition period, oviposition period, fecundity, incubation period and hatchability of eggs were recorded. The number of eggs laid per female was recorded daily, until the cessation of egg laying and the total number of eggs laid was worked out to obtain the fecundity. Number of larval instars, duration of each larval instar, total larval period, pupal period and total period of life cycle from egg to adult were also observed and the mean and standard deviation values were worked out. The life cycle was studied on both Jyothi and Aiswarya varieties of rice. The longevity of the beetles was recorded until death of the beetles and their mean and standard deviation values were worked out in both Jyothi and Aiswarya varieties.

#### RESULTS AND DISCUSSION

##### Site of oviposition, fecundity and oviposition period

The female beetle of *L. pygmaea* laid either single yellow oval shaped egg (Fig. 1) or in groups of 2, 3, 4 or 5 in a straight line on the upper or lower surface of mature or tender leaves of young rice plant thus indicating no preference for surface and age of rice leaf for oviposition. In the short duration variety Jyothi, it laid a mean of 16.8 eggs (Table 1) during an oviposition period of six days.

In the medium duration variety Aiswarya, a lower fecundity of 14.3 eggs was observed. A higher fecundity was thus observed in the short duration Jyothi than the medium duration Aiswarya. Egg laying was observed to be highest on the first day and thereafter it was reduced during the last fifth and sixth day on both the varieties. Eggs were seen glued to the dorsal or ventral leaf surface and no significant difference




FIGURE 1. Life stages of *Leptisma pygmaea*: a, eggs; b, c, d, e and f, 1st, 2nd, 3rd, 4th and 5th instar grub; g, Pupa; h, Adult beetle.

on the site of oviposition between rice varieties was observed and the eggs turned yellowish towards hatching.

In Maharashtra, Dalvi *et al.* (1985) observed that female blue beetle laid 38–66 eggs singly or in batches of 2, 3 or 4 in a straight line mostly on the lower surface. Very rarely eggs were observed on the upper surface of the leaves. Biological studies conducted in Gujarat (Patel and Shah, 1985) indicated that *L. pygmaea* laid an average of 43 and 58.78 eggs when temperature was above 30 °C and below 30 °C respectively showing a lower fecundity when temperature was above 30 °C. Dalvi *et al.* (1985) reported a higher fecundity of 43–58.78 eggs at a temperature range of 28–29 °C. The present study showed a lower fecundity for *L. pygmaea* which might be due to the higher temperature conditions prevailing in Kerala. An oviposition period of six days was observed in the present study, whereas in Maharashtra and Gujarat it was found to be 14 days (Patel and Shah, 1985; Dalvi *et al.*, 1985).

### Developmental stages

Eggs of *L. pygmaea* hatched into creamish white grubs within a mean period of 3.4 days (Table 1) in both Jyothi and Aiswarya thus indicating no difference between varieties. According to Patel and Shah (1985), the incubation period varied from 3.79

TABLE 1. Biology of *Leptispa pygmaea* on the rice varieties Jyothi and Aiswarya

Biological Parameters	Varieties of rice	
	Jyothi	Aiswarya
Mean number of eggs/Female	16.8 $\pm$ 1.23	14.3 $\pm$ 2.95
Mean egg period (Days)	3.4 $\pm$ 0.52	3.4 $\pm$ 0.52
Mean Larval period (Days)	8.2 $\pm$ 1.00	8.2 $\pm$ 0.40
Mean pupal period (Days)	3.2 $\pm$ 0.40	2.9 $\pm$ 0.57
Total life cycle (Days)	14.8 $\pm$ 2.15	13.8 $\pm$ 1.03
Pre-oviposition period (hours)	21.0 $\pm$ 4.10	20.3 $\pm$ 2.58
Adult male longevity (Days)	40.9 $\pm$ 12.17	36.7 $\pm$ 11.21
Adult female Longevity (Days)	24.9 $\pm$ 4.48	24.7 $\pm$ 3.50

$\pm$  indicates SD

TABLE 2. Duration of different larval stages of *L. pygmaea* on Jyothi and Aiswarya

Stage	Paddy variety	
	Jyothi	Aiswarya
Ist instar	2.2 $\pm$ 0.13	1.8 $\pm$ 0.42
IInd instar	1.6 $\pm$ 0.52	1.9 $\pm$ 0.32
IIIrd instar	1.3 $\pm$ 0.48	1.2 $\pm$ 0.42
IVth instar	1.3 $\pm$ 0.48	1.4 $\pm$ 0.52
Vth instar	1.8 $\pm$ 0.42	1.9 $\pm$ 0.74

$\pm$  indicates SD

to 7.16 days depending upon the temperature and the incubation period increased with decrease in temperature. They observed a constant level of 35 °C was lethal for hatching of eggs. The present result is in agreement with the findings of Patel and Patel (1970) who reported an incubation period of 4 to 5 days in *L. pygmaea*.

### Duration of different larval instars

In both Jyothi and Aiswarya, five instars were observed (Fig. 1) during the grub stage of rice blue beetle. The mean duration of 1st, 2nd, 3rd, 4th and 5th instars were found to be 2.2, 1.6, 1.3, 1.3 and 1.8 days respectively on Jyothi and the corresponding values on Aiswarya were 1.8, 1.9, 1.2, 1.4 and 1.9 days (Table 2). The average grub period (8.2 days) of *L. pygmaea* was found to be same in both varieties. Dalvi *et al.* (1985) observed that larva of *L. pygmaea* was fully developed in 12–14 days in Maharashtra. Patel and Shah (1985) also reported a higher larval period of 11.3 days for blue beetle in Gujarat. Five instars each with duration of 1–2 days were found in both Jyothi and Aiswarya. Patel and Shah (1985) observed only three instars for *L. pygmaea* in Gujarat.



### Pupal period

The grub pupated on the surface of the leaf as a brown coloured pupa (Fig. 1). The pupal period lasted for 3.2 and 2.9 days in Jyothi and Aiswarya respectively. Patel and Patel (1970) also reported that pupal period of *L. pygmaea* was about 4 days which is in agreement with the present finding. Patel and Shah (1985) observed that the pupal period were 4.14 days (at 25–28 °C) and 3.64 days (at 29–30 °C) indicating that pupal period was more in lower temperature. Dalvi *et al.* (1985) observed that the pupal period lasted for 4–5 days in Maharashtra.

### Total life cycle

*L. pygmaea* completed its life cycle within a mean period of 14.8 days in Jyothi while it was 13.8 days in Aiswarya. There was a difference of only one day in the mean duration of total life cycle of *L. pygmaea* on Jyothi and Aiswarya thus indicating the absence of marked difference in the duration of life cycle on short and medium duration rice varieties. In Maharashtra, Dalvi *et al.* (1985) reported that rice blue beetle completed its life cycle in 22–24 days. The shorter duration of life cycle in the present study might be due to the difference in weather conditions in the two regions.

### Pre-oviposition period

The mean pre-oviposition period was observed to be 21.0 and 20.3 h on Jyothi and Aiswarya respectively (Table 1). The mean pre-oviposition period in *L. pygmaea* was reported to be 4.9 days in Gujarat (Patel and Shah, 1985) and 2.6 days in Maharashtra (Dalvi *et al.*, 1985).

### Adult longevity

Longevity is important not only for its influence on demographics but also because it determines how long the pest is active and exerts a negative influence on the crop. The adult beetles are metallic blue coloured and black coloured on the ventral side of the body (Fig. 1). The longevity of adult varied with the sex of rice blue beetle. Male beetles lived longer than females. The male beetle had an average life span of 40.9 and 36.7 days in Jyothi and Aiswarya respectively (Table 1). The mean longevity of female beetle was 24.9 and 24.7 days in Jyothi and Aiswarya. Adult longevity was found to be higher on Jyothi than Aiswarya. This result is contrary to the earlier findings of Patel and Shah (1985) in Gujarat, where there was no marked difference in the longevity of male and female rice blue beetles. They found the adults lived a longer life of 19 days in March/April when maximum temperature fluctuated above 30 °C. During July–September, when maximum temperature was below 30 °C the adult beetles lived for 15–16 days. In Maharashtra, the adult beetles were found to have longevity of 18–35 days during kharif season (Dalvi *et al.*, 1985). The difference in the present results might be due to temperature and other prevailing rearing conditions.

## REFERENCES

- Dalvi, C. S., Dumbre, R. B. and Khanvilakr, V. C. (1985) Bionomics, seasonal incidence and off season biology of rice blue beetle. *Journal of Maharashtra Agricultural University* **10**(2): 185–187.
- Patel, G. A. and Patel, H. K. (1970) *Krishikit Vidya Part II*, Govt. of Gujarat, pp 57.
- Patel, C. B. and Shah, A. H. (1985) Biology of rice blue beetle, *Leptispa pygmaea* Baly. (Coleoptera: Hispinae). *Bulletin of Entomological Research* **26**(2): 120–128.
- Swamiappan, M., Rajaram, K. S. and Nair, V. C. (1990) Outbreak of blue beetle in India. *IRRI Newsletter* **15**(2): 16–17.

(Received 28 November 2007; accepted 15 February 2008)



## Insecticidal efficacy of *Chromolaena odorata* (compositae) on the coconut beetle, *Oryctes rhinoceros* (Linn.)

S. Leena<sup>1</sup>, B. T. Rayudu<sup>2</sup> and D. Muraleedharan<sup>\*3</sup>

<sup>1</sup>Central Plantation Crops Research Institute, Kasaragod 671124, India

<sup>2</sup>Krishi Vigyan Kendra, Central Plantation Crops Research Institute, Kasaragod 671124, India

<sup>3</sup>Centre for Arthropod Bioresources and Biotechnology, Department of Zoology, University of Kerala, Kariavattom Campus, Thiruvananthapuram 695581, India  
Email: leenakvk@gmail.com

**ABSTRACT:** Different doses of *Chromolaena odorata* plant powder were mixed with cowdung and fed to late third instar grubs of coconut rhinoceros beetle, *Oryctes rhinoceros* (L.). Mortality rates of larval and pupal stages as well as the emergence of deformed adults were recorded. Concentration of 15 and 20% (w/w) plant powder showed 80 and 90% impact respectively on the development of rhinoceros beetle. Topical supply of juvenile hormone analogue, methoprene @ 20 and 30  $\mu$ g levels to third instar grubs induced 86 and 92% developmental deformities similar to that observed with the supply of 15 and 20% plant powder. Similarity in the induction of morphological deformities in the emerging adults of *O. rhinoceros* by the two treatments indicated the JH mimicking endocrine influence exerted by this plant. These preliminary observations indicate the potential of this common weed for use against *O. rhinoceros*, by incorporating the plant material on the breeding sites of the pest. © 2008 Association for Advancement of Entomology

**KEYWORDS:** *Oryctes rhinoceros*, *Chromolaena odorata*, methoprene

### INTRODUCTION

Coconut rhinoceros beetle, *Oryctes rhinoceros* L. (Coleoptera: Scarabaeidae) is a serious and well known pest of coconut palm; the adult beetle bores into the unopened fronds and inflorescence of the palm. Adults breed on cow-dung or decaying organic matter on which the larvae feed. Treatment of breeding sites with insecticides is one of the key components of the IPM package for this pest. Application of insecticidal agent

\*Present address: Centre for Arthropod Bioresources and Biotechnology, Dept. of Zoology, University of Kerala, Thiruvananthapuram 695581.

at the breeding site, where the insect is relatively stationary, potentially would seem more efficient than other methods of control for the highly mobile adult (Campbell and Wright, 1976). However, this could be harmful to its natural predators and also can cause adverse effects on the environment. The present study was carried out to explore the possibility of eco-friendly management of this pest by incorporating plant product with insect control potential in its natural breeding sites. The plant tested is *Chromolaena odorata* (syn: *Eupatorium odoratum* L.). Commonly known as Siam weed, Christmas bush, etc., it is a noxious perennial weed in many parts of the world. It was introduced to India from tropical America during the Second World War and since then it has spread widely and has become a dominant weed of wastelands, roadsides and other exposed areas (Kushwaha *et al.*, 1981). Its medicinal properties are also well known.

Rajamma (1982) showed that the application of *C. odorata* leaves into the soil prior to planting of sweet potato reduces the weevil infestation significantly. The moult inhibiting action and formation of malformed adults as a result of application of *C. odorata* extract on *Dysdercus cingulatus* was reported earlier by Saradamma (1989). Prameela (1997) observed that the application of *C. odorata* extract resulted in the inhibition of normal ovarian development and oocyte maturation in the same insect. During fallows between cultivation, *Chromolaena* adds copious amounts of organic matter to the soil and may reduce the population of nematodes (M'Boob, 1991). It is also an useful mulch for row crops (Swennen and Wilson, 1984). Basin application of *Chromolaena* leaves prior to planting of vegetable seedlings is recommended and followed by vegetable growers of Kerala. In pepper, evaluation of leaf and seed extracts of *Strychnos nuxvomica* and *Chromolaena odorata* were promising indicating their potential for utilizing them in IPM schedules (Devasahayam, 2005).

Keeping the above in view, the present study was undertaken. Studies were also conducted on the effect of synthetic juvenile hormone mimic, methoprene on the development of rhinoceros beetle for comparison of observed symptoms.

#### MATERIALS AND METHODS

Late third instar *O. rhinoceros* (L.) grubs were used for the experiment. Grubs of different stages of growth were collected from local manure pits, vermi-compost units and felled oil-palm residues and were maintained on cow-dung in the laboratory under controlled conditions. Tender stems and leaves of *C. odorata* were collected from Kariavattom campus and nearby areas, shade-dried and pounded to a coarse powder. This powder @ 5, 10, 15 and 20% (w/w) was thoroughly mixed with sun-dried and powdered cow-dung. Cow-dung without leaf powder served as control. Late third instar grubs with a mean weight of 11.2 g were introduced individually into small plastic containers with 100 g of cow-dung-plant mixture. Water was sprinkled over the feed to provide adequate moisture. The containers were covered with plastic lids with holes. The feed was changed once a week. There were 10 grubs per treatment and the experiment was replicated five times. Observations were made on the development of the larvae and deformities, if any, at three days intervals.

TABLE 1. Effect of *Chromolaena odorata* leaf powder on the development of *Oryctes rhinoceros*

Dosage (%plant powder)	Initial no. of larvae	Larval mortality	Pupal mortality	Total mortality	Deformed adults	Deformity score	Total mortality+ Deformed adults
05	10	1.4	0.4	1.8	1.4	0-1	3.2
10	10	2.6	0.6	3.2	3.2	1-2	6.4
15	10	3.6	1.2	4.8	3.2	2-3	8.0
20	10	6.0	1.2	7.2	1.8	3-4	9.0
Nil (control)	10	0.2	0.0	0.2	0.0	-	0.2

Values represent the mean of 5 replications with 10 larvae each.

General mean: 1.787; S.E: 0.560; CV(%): 21.330

Treatments (T) CD: 0.409; Conditions (C i.e., Developmental stages of the insect) CD: 0.317; T X C CD: 0.709.

In another experiment, the effect of juvenile hormone analogue, methoprene on the development of third instar *O. rhinoceros* larva was studied. Methoprene (ZR 515, gift from Dr. Govindan Bhaskaran, A&M University, Texas, USA) was dissolved in acetone so that 1  $\mu$ l contained 1  $\mu$ g methoprene and the desired dose (10  $\mu$ g, 20  $\mu$ g and 30  $\mu$ g) was applied to different animal groups of ten each in five replications. The hormone was applied topically on the ventral abdominal segment of third instar larvae. Two sets of control were maintained; one set received only acetone in the same dosages and the other received no treatment at all. All the sets of larvae were maintained on cow-dung as described earlier. The treatments were replicated five times and observations recorded as above.

## RESULTS

When third instar *O. rhinoceros* larvae were grown on cow-dung mixed with *C. odorata* leaf powder, the first larval mortality was recorded on the sixth day of treatment. Larval mortality was 60% in 20% leaf powder medium, followed by 36% in 15% leaf powder, 26% in 10% leaf powder and 14% in 5% leaf powder (Table 1). Only 2% larval mortality was observed in untreated cow-dung. In case of pupal mortality, 12% each in 20% and 15% leaf powder was recorded followed by 6% and 4% in 10% and 5% leaf powder, respectively. The percentage of abnormal adults was 32 in 15% and 10% leaf powder, 18 in 20% leaf powder and 14 in 5% leaf powder. Pupal mortality and abnormal adults were not found in control. In 20% leaf powder medium, 90% of the insects were affected either by mortality of larva or pupa or emergence of abnormal adults. The percentage of insects thus affected was 80 in 15% leaf powder, 64 in 10% leaf powder and 32 in 5% leaf powder.

Pupal-adult intermediates were observed in treatments, which were characterized by adult head, pupal wing buds and retention of pupal exuviae. They lived only up to five

TABLE 2. Effect of methoprene on the development of *Oryctes rhinoceros*

Dosage	Initial no. of larvae	Larval mortality	Pupal mortality	Total mortality	Deformed adults	Deformity score	Total mortality+ Deformed adults
10	10	1.2	0.2	1.4	4.8	1-2	6.2
20	10	2.0	0.4	2.4	6.2	3-4	8.6
30	10	3.4	0.8	4.2	5.0	2-3	9.2
Acetone							
30	10	0.0	0.0	0.0	0.0	—	0.0
Nil	10	0.0	0.0	0.0	0.0	—	0.0

Values represent the mean of 5 replications with 10 larvae each.

General mean: 1.573; S.E: 0.510; CV(%): 22.439

Treatments (T) CD: 0.373; Conditions (C) (i.e. Developmental stages of the insect) CD: 0.289; T x C CD: 0.647

days whereas those adults with badly deformed wings survived up to nine days. The emerged adults other than malformed were normal looking but with pale, leathery, unchitinized wings not covering the abdomen fully. Around 80% of them had longevity of 24–30 days. From control larvae, normal adults emerged and lived beyond 95 days. More than 80% of emerged adults were females and abnormal morphological characters were manifested more in females. The pygidium of normal looking and deformed adult females under plant product treatment was not densely packed with reddish-brown sensory hairs whereas it was bushy in control adult females. This may be due to some abnormalities in the reproductive system of treated larvae.

When treated with methoprene, 34% larval mortality was recorded at 30  $\mu$ g dose, followed by 20% at 20  $\mu$ g and 12% at 10  $\mu$ g (Table 2). Pupal mortality was 8%, 4% and 2% respectively at the above doses. Highest abnormal adults (62%) was recorded at 20  $\mu$ g methoprene followed by 50% at 30  $\mu$ g and 48% at 10  $\mu$ g. Treatment with 30  $\mu$ g methoprene affected 92% of the insects either by mortality of larva or pupa or production of abnormal adults; 20  $\mu$ g methoprene affected 86% of the insects and 10  $\mu$ g methoprene affected 62% of the insects. The morphological deformities observed in adults emerged out of this treatment was similar to that produced by exposure to *C. odorata* leaf powder. No larval or pupal mortality or abnormal adults were found in the two controls.

## DISCUSSION

From the present results, it is inferred that *Chromolaena odorata* has exerted JH like activity during the metamorphosis of this insect as evidenced through the emergence of adults with varying external deformities, occurrence of pupal-adult intermediates and premature death of adults inside the pupal case. The chemical properties of juvenoids usually allowed them to cross the cuticle and act directly on the insect as a result of contact. Since ingested JH like substances are likely to be digested before absorption,

the effect observed here seems to be due to external skin contact of larvae with the plant material.

Growth regulatory activity including larval and pupal mortality as well as deformation of adults in *Oryctes rhinoceros* was reported with *Clerodendron infortunatum* (Chandrika and Nair, 2000). Further, the results of the present study agree with the findings of Ponnammma (2003) that crushed leaves of *C. odorata*, when mixed with sterilized cow dung and fed to third instar grubs of *O. rhinoceros*, caused abnormalities and prevented proper development of beetles. The insecticidal and insect growth regulatory activity exerted by organic extracts from the leaves of *Vitex mollis* was recently reported by Rodriguez-Lopez *et al.* (2007). Plant kingdom offers an excellent source of secondary metabolites which affect the behaviour, physiology, growth, reproduction and development of insects. Many of them exert repellence effect, feeding deterrence and several other influences. Some affect the hormonal balance of the insects while some others are highly toxic and affect normal physiological processes. Many juvenoids have been isolated from plants and the possibility of using these phytojuvenoids in disrupting the insect developmental processes forms an innovative and effective control strategy for the management of insect pests. The use of these growth regulating compounds would be more appropriate, when the insect is a pest during its adult stage and the larvae develop in a restricted habitat (Elzinga, 2000). The effects produced by *C. odorata* leaf powder and the JHa methoprene on *O. rhinoceros* showed resemblances in their morphological abnormalities indicating the presence of juvenile hormone mimics in the tested plant.

The search and use of safer plant-derived products are increasing now as a viable component of Integrated Pest Management (IPM). Results of the present study suggest the possibility of incorporating *C. odorata*, a common weed plant available in plenty in coconut orchards, in the breeding sites of *O. rhinoceros*, as an ecologically safer, human hazard free, farmer-friendly component among the existing IPM package developed for the management of *O. rhinoceros*.

#### ACKNOWLEDGEMENTS

The first author thanks the Director, Central Plantation Crops Research Institute, ICAR, Kasaragod for grant of study leave to undertake research work leading to Ph.D degree at Kerala University, and Professor & Head, Department of Zoology, Kerala University, Trivandrum for extending facilities.

#### REFERENCES

- Campbell, J. B. and Wright, J. E. (1976) Field evaluation of insect growth regulators, insecticides and bacterial agent for Stable fly control in feedlot breeding areas. *Journal of Economic Entomology* **69**(5): 566–568.
- Chandrika, Mohan and Nair, C. P. R. (2000) Effect of *Clerodendron infortunatum* on grubs of coconut rhinoceros beetle, *Oryctes rhinoceros* L. In: *Recent Advances in Plantation Crops Research*, Muraleedharan, N. and Raj Kumar, R. (Eds). 297–299.
- Devasahayam, S. (2005) Potential use of plant products for the management of insect pests of spices. In: *Paper presented in National Seminar on Insect Growth Regulators and*

- Natural Products in Insect Pest Management*, organized by P.G and Research Department of Zoology, St. Joseph's College, Devagiri, Calicut.
- Elzinga, R. J. (2000) Integrated pest management. In: *Fundamentals of Entomology-Fifth Edition*, Published by Charles Stewart, 314-334.
- Kushwaha, S. P. S., Ramakrishnan, P. S. and Tripathi, R. S. (1981) Population dynamics of *Eupatorium odoratum* in successional environments following slash and burn agriculture. *Journal of Applied Ecology* **18**: 529-535.
- M'Boob, S. S. (1991) Preliminary results of a survey and assessment of *Chromolaena odorata* (Siam weed) in Africa. *Biotropica Special Pub.* **44**: 51-55.
- Ponnamma, K. N. (2003) Integrated Pest Management practices for Rhinoceros beetles and Red Palm Weevil of Oil palm, National Agriculture Technology Project, Final Report
- Prameela, M. (1997) Bio-insecticides in the physiology of red cotton bug, *Dysdercus cingulatus* (Heteroptera; Pyrrhocoridae) *Ph.D. Thesis*, Kerala University, Trivandrum.
- Rajamma, P. (1982) Effect of some organic materials on the control of sweet potato weevil *Cylas formicarius* Fab. *Journal of Root Crops* **8(1-2)**: 64-65. CTCRI, Trivandrum
- Rodriguez-Lopez, V., Figueroa-Suarez, M. Z., Rodriguez, T. and Aranda, E. (2007) Insecticidal activity of *Vitex mollis*. *Fitoterapia* **78(1)**: 37-39.
- Saradamma, K. (1989) Biological activity of different plant extracts with particular reference to their insecticidal, hormonal and antifeeding actions *Ph.D. Thesis*, KAU, Thrissur.
- Swennen, R. and Wilson, G. F. (1984) *In-situ* mulch production for plantain. *Banana Newsletter* **7**: 20-22.

(Received 4 January 2008; accepted 15 February 2008)





## Population dynamics, species composition and feeding site preference of rice black bug (hemiptera: pentatomidae) infesting rice

**P. Anandhi\*, M. A. K. Pillai and Savita Varma**

*Department of Agricultural Entomology, AC & RI, Killikulam, Tamil Nadu  
Agricultural University, Tamil Nadu, India*

**ABSTRACT:** The incidence of rice black bug in different periods in rice fields was monitored in South Tamil Nadu. Maximum nymphs and adults were recorded during January. This coincided with 97th day after transplanting (DAT) followed by 87 and 45 DAT. Two species of black bug were seen in the population, *Scotinophara coaractata* (Fab.) and *S. lurida* (Burm.), the latter being very scarce (2.5 to 1.0 % of total catch) in all the months. The nymphs and adults preferred the leaf node for feeding. © 2008 Association for Advancement of Entomology

**KEYWORDS:** rice black bug, *Scotinophara coaractata*, stink bug

### INTRODUCTION

Stink bugs of the genus *Scotinophara* are common pests of rice in several Asian countries (Miyamota *et al.*, 1983). Of the six *Scotinophara* sp. present on rice in the Phillipines, only *S. coaractata* (F.) was reported as a serious pest (Barrion *et al.*, 1982). Earlier reports suggested that it was an introduced pest (Miyamota *et al.*, 1983) but it may be that its pest status has only recently been recognized. Though it is reportedly an introduced pest in many countries its entry into India lacks evidence. In Tamil Nadu introduced pest in many countries its entry into India lacks evidence. In Tamil Nadu sporadic but severe outbreaks of the pest was reported by Uthamasamy and Marriappan (1985) and Subramanian *et al.* (1986). Rice black bug hitherto considered as a minor pest has become a menace during the past five years.

### MATERIALS AND METHODS

The present investigation was undertaken to study the population dynamics, species complex and the host preference of black bugs infesting rice at Agricultural College

\*Present address: Department of Plant Protection Allahabad Agricultural Institute-Deemed University, Allahabad 7, UP

and Research Institute, Killikulam, Tamil Nadu. For studying the population dynamics a field experiment was conducted in the central farm of the college in Pishanam season planting ASD-16 variety and following recommended agronomic practices. The population was assessed by counting the number of nymphs and adults visually from 40 hills at random commencing from 12 days after transplanting at monthly intervals. The affected hills were removed, placed in plastic bags and transported to the laboratory and the nymphs and adults were counted and recorded. Each represented a sampling unit. Sampling was carried out over a ten months period.

Population abundance of black bug during one cropping season was also assessed in IR-20 variety in the field. The total population of nymphs and adults were counted in 10 hills at random from 27–97 days after transplanting at 10 day intervals. The identification of species complex was made through light trap collection. The light traps were installed in the field and the trap was fitted with incandescent bulb (100W). Daily catches collected between 7 PM to 6 AM were counted. Adult pentatomid bugs of *Scotinophara* sp. collected from the light traps were separated, sorted out and identified by taxonomic characters (Barrion *et al.*, 1982). Predominant species in every month were recorded.

In a net house study the preferred feeding sites of black bug was observed. Bugs were released on 30–35 days old ADT-36 (a rice variety highly susceptible to black bug) at panicle initiation stage and covered with Mylar cages which enclosed 7 adults/plant with suitable control (plant with no bug). Treatments were replicated thrice. The insect feeding at nodal, stem, leaf and panicle stages were assessed.

## RESULTS AND DISCUSSION

Data on the incidence of black bug in different periods in field condition are shown in Table 1. Weekly in situ counts of the nymphs and adults from 40 hills at random revealed maximum population in the month of January (220 numbers) and beyond that the incidence gradually decreased. Population of nymphs and adults showed the same trend. Ferer and Shepard (1987) studied the light trap catches of black bug and recorded the peak activities in mid August and mid April and June. The effect of growth stage of the plants on the population of black bug is presented in Table 2. The population significantly varied among the age of the crop. The mean maximum population was recorded at 97 DAT (162 numbers) followed by 87 DAT (141 numbers) and 47 DAT (113 numbers). This is in agreement with the earlier reports of Abdul Latif *et al.* (1982) and Saroja *et al.* (1993).

Table 3 shows the species of black bug in different periods. The data clearly establish the dominance of *S. coaractata* throughout the cropping season compared to *S. lurida*. Peak population was recorded in March, May, June and September (100%) and smaller peaks were evident in February (94.6%). Saroja (1991) recognized that *S. coaractata* was the predominant species and accounted for 97% incidence in Chengalpattu district of Tamil Nadu. Within the district level survey occurrence of the two species showed variations and *S. coaractata* was the most common species.

TABLE 1. Seasonal incidence of black bug in different periods in field condition

Months	Population No./40 hills		
	Nymph	Adult	Total
Dec	50 (7.07) <sup>f</sup>	112 (10.58) <sup>d</sup>	162 (17.65) <sup>e</sup>
Jan	80 (8.95) <sup>b</sup>	140 (11.83) <sup>a</sup>	220 (20.78) <sup>a</sup>
Feb	67 (8.15) <sup>d</sup>	126 (11.23) <sup>b</sup>	193 (19.42) <sup>d</sup>
Mar	40 (6.33) <sup>h</sup>	93 (9.64) <sup>e</sup>	133 (15.97) <sup>f</sup>
Apr	27 (5.26) <sup>e</sup>	46 (6.78) <sup>h</sup>	73 (12.04) <sup>h</sup>
May	15 (3.87) <sup>j</sup>	32 (5.66) <sup>i</sup>	47 (9.53) <sup>j</sup>
June	47 (6.86) <sup>g</sup>	22 (4.69) <sup>j</sup>	69 (11.55) <sup>i</sup>
July	136 (11.66) <sup>a</sup>	71 (8.43) <sup>f</sup>	207 (20.09) <sup>b</sup>
Aug	59 (7.68) <sup>i</sup>	58 (7.62) <sup>g</sup>	117 (15.30) <sup>g</sup>
Sep	78 (8.83) <sup>c</sup>	123 (11.09) <sup>c</sup>	201 (19.92) <sup>c</sup>

Values represent the mean of three replications. In a column, means followed by the common letter are not significantly different ( $P = 0.05$ ) by LSD; DAT - Days after transplanting.

TABLE 2. Population of black bug at different periods of crop growth observed in field condition

Days after transplanting	Population No./10 hills		
	Nymph	Adult	Total
27	0.5 (0.71) <sup>a</sup>	8 (2.83) <sup>a</sup>	8.5 (3.54) <sup>a</sup>
37	23 (4.79) <sup>b</sup>	58 (7.62) <sup>d</sup>	81 (12.47) <sup>b</sup>
47	54 (7.35) <sup>d</sup>	59 (7.68) <sup>d</sup>	113 (15.03) <sup>e</sup>
57	52 (7.21) <sup>c</sup>	43 (6.56) <sup>c</sup>	95 (13.77) <sup>d</sup>
67	58 (7.62) <sup>e</sup>	35 (5.92) <sup>b</sup>	93 (13.54) <sup>c</sup>
87	69 (8.31) <sup>f</sup>	72 (8.49) <sup>e</sup>	141 (16.80) <sup>f</sup>
97	52 (7.21) <sup>c</sup>	110 (10.49) <sup>f</sup>	162 (17.70) <sup>g</sup>

Values represent the mean of three replications. In a column, means followed by the common letter are not significantly different ( $P = 0.05$ ) by LSD.

TABLE 3. Species complex of *Scotinophara* in different periods obtained from light trap catch

Months	<i>S. coaractata</i>		<i>S. lurida</i>	
	No.	Percentage in total catch	No.	Percentage in total catch
Dec	156	97.5	4	2.5
Jan	289	98.0	6	2.0
Feb	35	94.6	6	2.0
Mar	10	100.0	0	—
Apr	192	99.0	2	1.0
May	5	100.0	0	—
June	9	100.0	0	—
July	85	97.7	2	2.3
Aug	39	97.5	1	2.5
Sep	7	100.0	0	—
Mean	82.7	98.4	17	1.57

TABLE 4. Preferred feeding sites of black bug

Plant part fed	Insect feeding in %	
	Nymphs	Adults
Node	73 <sup>a</sup>	88 <sup>a</sup>
Stem	68 <sup>a</sup>	73 <sup>b</sup>
Leaf	43 <sup>b</sup>	8 <sup>d</sup>
Panicle	28 <sup>c</sup>	18 <sup>c</sup>

Values represent the mean of three replications. In a column, means followed by the common letter are not significantly different ( $P = 0.05$ ) by LSD.

The preference of feeding sites of the black bug is shown in Table 4. Among the different plant parts, nodal portion was most preferred by the adults and nymphs (88 and 73%) followed by stem (73 and 68%) and leaf petiole (8 and 43). This is in consonance with the report of Saroja *et al.* (1993).

## REFERENCES

- Abdul Latif, A. Z., Chang, P. M. and Nik Mohamed, Noor (1982) The incidence and chemical control of the Malayan black bug *Scotinophara coaractata* in west Malaysia. *Proc. Pad. Workshop. Illus. Bombong Lima, Mardi* 276–289.
- Barrion, A. T., Mochida, O., Litsinger, J. A. and Delacruz, N. (1982) The Malayan black bug, *S. coaractata* (F.) (Hemiptera: Pentatomidae) a new rice pest in the Philippines. *Int. rice Res. Newsl* **7**(6): 6–7.
- Ferer, E. R. and Shepard, B. M. (1987) Sampling Malayan black bugs (Heteroptera: Pentatomidae) in rice. *Environ. Entomol.* **16**: 259–263.
- Miyamota, S., Torres, N. A., Habibuddin, B. H., Montri, R., Fujimura, T., Thieng, P. and Mochida, O. (1983) Proceedings emerging problems of pests and Diseases black bug in south east Asia. In: *Int. Rice Res. Conference*, Los Banos, Phillipines.
- Saroja, R. (1991) Black bug incidence in rice. In: *Paper Presented in the 8th State Level Plant Protection Workers Meet. Nov. 1991*, TNAU, Coimbatore, India.
- Saroja, R., Subramanian, M. and Abdul Kareem, A. (1993) *Rice Black Bug*, Rice Research Station, Tamil Nadu Agric. Univ., Tirur.
- Subramanian, A., Murugesan, S., Rajendran, R. and Sundara Babu, P. C. (1986) Occurrence and control of rice black bug at coimbatore. *Int. Rice Res. Newsl.* **11**(3): 24.
- Uthamasamy, S. and Marriappan, V. (1985) Occurrence of black bug in Tamil Nadu. *Int. Rice Res. Newsl.* **10**(2): 15.

(Received 29 June 2007; accepted 15 February 2008)





## **Revival and recharacterization of genus *Thyrgorina* walker (Lepidoptera: Arctiidae: Arctiinae) and taxonomic studies on four Indian species of this genus from Western Ghats of India**

**Jagbir Singh Kirti and Navneet Singh Gill**

*Department of Zoology, Punjabi University, Patiala 147002, Punjab, India*

**ABSTRACT:** Genus *Thyrgorina* Walker has been revived from the synonymy of *Diacrisia* Hübner and recharacterized in detail by including external male and female genitalic structure of its type species. External genitalic structure of Indian species i.e. *Thyrgorina indica* (Guer), *T. angularis* (Strand) and *T. exima* (Swinhoe) have been studied and illustrated for the first time besides reporting of a new species viz. *T. coorgensis* sp.nov. from Western Ghats of India. The description of new species is discussed in detail. © 2008 Association for Advancement of Entomology

**KEYWORDS:** Arctiinae, Lepidoptera, *Thyrgorina*, revalidation, new species

### **INTRODUCTION**

Genus *Thyrgorina* was established by Walker in 1864 with *indica* Guer as the type species with range from Himalayas to Nilgiris in India, and Myanmar. Hampson in 1894 described as many as 14 species under genus *Thyrgorina* Walker from different areas of then British India. Hampson (1901) synonymised this genus under *Diacrisia* Hübner and the same nomenclature was followed by Strand in 1922. Koda (1988) while giving the generic classification of Arctiinae of the Palearctic and Oriental regions based on the male and female genitalia shifted some of the species previously placed under genus *Thyrgorina* Walker to *Thanatarctia* Butler suggesting new combinations.

During the present studies, as many as four species i.e., *indica* Guer, *angularis* Strand, *eximia* Swinhoe and *malshejensis* sp. nov. were collected and studied in detail. The critical examination of the type species of genus *Thyrgorina* Walker i.e., *indica* Guer reveals that it is generically distinct from type species of genus *Diacrisia* Hübner. The male and female genitalic structures of remaining three species have also been studied in detail and these are completely congeneric with the type species *indica* Guer. In this light, genus *Thyrgorina* Walker has been removed from the synonymy

of genus *Diacrisia* Hübner and is being revived and recharacterized by incorporating male and female genitalic characters in its diagnosis. A key to the four studied species is given below. One of the species could not be identified from the relevant literature or by comparison from different national museums and Natural History Museum, London. Therefore, it is being proposed as new species, *malshejensis* sp. nov. here.

#### MATERIALS AND METHODS

The material for the present studies i.e. the adult moths of sub-family Arctiinae were collected at night using fluorescent light from different localities of Western Ghats of India. The moths thus collected were killed with ethyl acetate in the killing bottle. The freshly killed specimens were pinned and stretched on adjustable wooden stretching box. The pinned specimens were dried for 2–3 days in the improvised drying chambers fitted with electric bulb. The properly dried specimens were then preserved in air tight wooden boxes, containing naphthalene balls as fumigant. The method proposed by Common (1970) and advocated by Zimmerman (1978) has been followed for the preparation of permanent slides of fore and hindwings. For the study of external male genitalia, the entire abdomen of preserved moths was removed, as cutting the last few segments often damages the constituent parts of male and female genitalia (Robinson, 1976). The detached abdomen was put in 10% KOH for 12–14 h to soften the chitin and to dissolve away the muscles and other soft parts. The KOH treated material was washed in distilled water and residual traces of KOH were later removed using 1% glacial acetic acid. The abdomen was dissected in 50% alcohol for taking out the genitalia and the adhering unwanted material was cleared off in the subsequent grades. After proper dehydration, the material was cleared in clove and preserved in alcohol-glycerol. The diagrams were drawn with the help of graph eye piece fitted in the zoom binocular.

#### RESULTS AND DISCUSSION

##### **Genus Thyrgorina Walker gen. rev.**

Walker, 1964, *List. Spec. Lepid. Ins. Coll. Br. Mus.*, **31**: 37.

##### **Type species** *Arctia indica* Guer.

**Distribution** India: Himalayas, Mumbai, Nilgiri Hills (Tamil Nadu); Myanmar.

##### *Diagnosis:*

Labial palpi porrectly downward. Antennae bipectinate in males, serrate in females. Forewing with vein  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  stalked;  $M_1$  from upper angle of cell;  $M_2$  and  $M_3$  from lower angle of cell. Hindwing with vein  $Sc + R_1$  originating towards base of cell;  $M_2$  and  $M_3$  from or near lower angle of cell. Hind tibia with two pair of spurs. Male genitalia with uncus long, broad at base and narrowing towards tip; acrotergite weakly developed; fenestrula triangular; valvae simple with costa narrow and sacculus



well differentiated, ampula not distinct; aedeagus long; vesica membranous with well formed patch of sclerotized spines; ductus ejaculatorius entering subapically. Female genitalia with corpus bursae membranous, signum absent.

***Thyrgorina indica* (Guer) comb. rev. (Fig. 1)**

*Arctia indica* Guer, 1843, *Deless, Souv. Inde.*, 2: 9.3

*Thyrgorina spilosomata* (Walker, 1864), *Cat. Lep. Het.*, 31: 318.

*Thyrgorina indica* (Guer, 1843) Hampson, 1894, *Fauna Br. Ind. Moths*, 2: 12

*Diacrisia indica* (Guer, 1843) Hampson, 1901, *Cat. Lep. Het.*, 3: 265

*Male genitalia*

Uncus strongly built, beak-like, well sclerotized, setosed with small seate, broad at base, tip pointed; acrotergite weakly developed; fenestrula triangular; tegumen as long as uncus, u-shaped; vinculum slightly shorter than tegumen, weakly sclerotized, v-shaped; saccus present. Valvae with costa and sacculus differentiated; harpe + ampula simple; cucullus produced to a snout, tip blunt; valvula a small outgrowth on lateral side. Juxta oval with a protrusion at distal end; aedeagus long and narrow; vesica membranous with series of sclerotized spines; ductus ejaculatorius entering subapically.

*Female genitalia*

Corpus bursae funnel shaped, broad at posterior region and cylindrical towards anterior region, membranous, signum absent; ductus bursae moderately long, broad at both ends and narrowing towards central region, weakly sclerotized; ductus seminalis entering at junction of corpus bursae and ductus bursae; anterior apophyses shorter than posterior apophyses; papilla analis fringed with short setae.

**Wing span** Male 32–34 mm; female 40–44 mm.

**Material examined**

Gujarat:	Saputara (970 m), 29. ix.05 – 2♂♂
Maharashtra:	Matheran (690 m), 5.x.05 - 9♂♂ 1♀; Mahabaleshwar (1320 m), 8.x.05 – 1♂; Malshej Ghat (690 m), 1.x.05 1♂.
Tamil Nadu:	Coonoor (1880 m), 2.x.03 - 1♂
Kerala:	Devikulam (1620 m), 13.ix.04 – 1♂, 14.ix.04 - 4♂♂; Vallakadav (780 m), 11.ix.04 – 3♂♂ 1♀.

**Old distribution** India: Mumbai and Nilgiri Hills (Tamil Nadu)

*Remarks*

As mentioned under the remarks of genus, the original status of genus *Thyrgorina* Walker has been restored in the present study and the external male and female genitalia of its type species i.e. *indica* Guer are discussed and illustrated here in detail.




FIGURE 1. *Thyrgorina indica* (Guer) comb. rev. (A) Forewing; (B) Hindwing; (C) Male genitalia; (D) Aedeagus; (E) Valva (right); (F) Valva (left); (G) Uncus with Tegumen (dorsal view); (H) Uncus with Tegumen (lateral view); (I) Juxta (enlarged); (J) Female genitalia.

***Thyrgorina angularis* (Strand) stat. nov. (Fig. 2)**

*Diacrisia indica* Guer ab. *angularis* Strand; Strand 1919, *Arctidae: Arctiinae*.  
*Lep. Cat.* 22: 182.

**Head** Frons and vertex white. Antennae bipectinate in males, serrate in females; scape and pedicel covered with white scales; shaft and branches brown with slight suffusion of white scales. Eyes fuscus grey, densely spotted with black. Labial palpi porrectly downward; underside covered with ochreous scales; lateral side black; third segment black and bent downward.

**Thorax** White; tegula furnished with white hair like scales; collar white having suffusion of yellow scales.

**Forewing** Ground colour white; base of the costa black brown; a spot at end of cell; veins paler; underside same with postmedial series of three spots; fringe white; vein  $R_1$  originating from cell;  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_5$  stalked;  $M_1$  from upper angle of cell;  $M_2$  and  $M_3$  from lower angle;  $Cu_1$  near lower angle;  $Cu_2$  from middle of cell.

**Hindwing** Ground colour white; a black spot at end of cell; submarginal black spots under vein  $Cu_2$ ; a black spot at vein  $M_2$ ; underside same; fringe white; vein  $Sc + R_1$  originating near base of cell;  $Rs$  and  $M_1$  from upper angle of cell;  $M_2$  and  $M_3$  from lower angle;  $Cu_1$  very near to lower angle;  $Cu_2$  from middle of cell. Legs covered with white scales; outer surface of tibia and tarsus black brown; hind tibia with two pair of spurs.

**Abdomen** Yellow with first segment white; dorsal, lateral and sublateral series of black spots; underside and tuft white.

*Male genitalia*

Uncus broad at base and narrowing towards tip, setosed with small setae, well sclerotized, tip pointed and curved; acrotergite poorly developed; fenestrula triangular; tegumen longer than uncus, u-shaped; vinculum shorter than tegumen, sclerotized, u-shaped, saccus developed. Valvae simple with costa weakly sclerotized; sacculus well sclerotized, produced to an outgrowth towards distal end; harpe+ampulla simple; cucullus simple; valvula with a small outgrowth. Transtilla weakly sclerotized; juxta almost rectangular with a protrusion towards distal end and a notch at proximal end; aedeagus moderately long and broad; vesica membranous with a patch of small sclerotized spines; ductus ejaculatorius entering subapically.

*Female genitalia*

Corpus bursae oval, membranous, signum absent; ductus bursae moderately long and broad, well sclerotized; ductus seminalis entering at junction of ductus bursae and corpus bursae; anterior apophyses shorter than posterior apophyses; papilla analis setosed with long setae.

**Wing span** Male 32 mm; female 40–42 mm.




FIGURE 2. *Thyrgorina angularis* (Strand) stat. nov. (A) Forewing; (B) Hindwing; (C) Male genitalia; (D) Aedeagus; (E) Valva (right); (F) Valva (left); (G) Uncus with Tegumen (dorsal view); (H) Uncus with Tegumen (lateral view); (I) Juxta (enlarged); (J) Female genitalia.

**Material examined**

- Gujarat: Ahwa (520 m), 28.ix.05 – 1♂; Saputara (970 m), 29.ix.05- 2♂♂; 3♀♀.
- Maharashtra: Mahabalehwar (1320 m), 8.x.05 – 1♂, 9.x.05 – 1♂, Malshej Ghat (690 m), 1.x.05 – 1♂.
- Kerala: Devikulam (1620 m), 13.ix.04 – 1♀, 14.ix.04 – 1♂.

**Old distribution** India: Mumbai, Nilgiri Hills (Tamil Nadu) and Coimbatore.

**Remarks**

While describing species *indica* Guer, Hampson (1901) discussed its representatives with spots of forewing reduced to one at upper angle of cell as a distinct 'form' under *Diacrisia indica* Guer. Strand (1919) named this 'form' as *angularis* Strand. The detailed study of the species under reference reveals that, besides external morphological characters it can be separated from *indica* Guer on the basis of external genitalic structures like shape of juxta, arrangement of spines in vesica of male genitalia and distinct shape of corpus bursae of female genitalia. Therefore, the 'form'; described by Hampson and Strand i.e. *angularis* Strand has been upgraded here by giving it its specific status.

**THYRGORINA MALSHEJENSIS SP. NOV. (FIG. 3)****Head**

Frons and vertex ochreous. Antennae bipectinate in males, serrate in females; scape, pedicel and shaft covered with buff scales; branches brown. Eyes yellowish green, densely spotted with black. Labial palpi porrect downward; laden with ochreous scales; tip of third segment black.

**Thorax** Ochreous, collar and tegula covered with ochreous bordered fuscus, spots; pectus ochreous.

**Forewing**

Ground colour ochreous; a basal brown patch; diffused subbasal, antemedial, medial and postmedial brown bands; underside same; fringe ochreous; vein R<sub>1</sub> from cell; R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> from a common stalk, M<sub>1</sub> near upper angle of cell; M<sub>2</sub> and M<sub>3</sub> from lower angle of cell; Cu<sub>1</sub> near lower angle; Cu<sub>2</sub> from middle of cell.

**Hindwing**

Ground colour ochreous; discal cell, vein IA and median nervure suffused with brown scales; more or less complete post medial brown bands; underside same; fringe ochreous; vein Sc+R<sub>1</sub> originating before middle of cell, Rs and M<sub>1</sub> on a short stalk; M<sub>2</sub> and M<sub>3</sub> from lower angle; Cu<sub>1</sub> near angle of cell; Cu<sub>2</sub> from middle of cell. Legs ochreous with slight suffusion of fuscus scales; hind tibia with two pairs of spurs.

**Abdomen** Orange yellow; dorsal and lateral series of black spots; underside covered with buff scales.




FIGURE 3. *Thyrgorina malshejensis* sp. nov. (A) Forewing; (B) Hindwing; (C) Male genitalia; (D) Aedeagus; (E) Valva (right); (F) Valva (left); (G) Uncus with Tegumen (dorsal view); (H) Uncus with Tegumen (lateral view); (I) Juxta (enlarged).

*Male genitalia*

Uncus long, broad at base and gradually narrowing towards tip, slightly wavy at ventroproximal region, setosed with small setae, tip minutely blunt; acrotergite present; fenestrula triangular; tegumen longer than uncus, u-shaped; vinculum shorter than tegument, well sclerotized, deep u-shaped, saccus weakly developed. Valvae with costa narrow and linear, weakly sclerotized; sacculus well differentiated, folded, produced to an outgrowth towards proximal end; harpe + ampulla simple plate like; cucullus with a small outgrowth, produced to narrow snout, tip rounded; valvula with small setae, well sclerotized. Transtilla sclerotized; juxta pot shaped; aedeagus long and broad, almost straight; vesica membranous with a single patch of well formed large sclerotized spines; ductus ejaculatorius entering laterally.

**Wing span** Male 32 mm.

**Material examined**

Holotype:	Maharashtra:	Malshej Ghat (690 m), 2.x.05 -1♂
Paratype:	Maharashtra:	Malshej Ghat (690 m), 2.x.05 -1♂

**Remarks**

The species under reference is closely allied to *Thyrgorina flavens* Moore as far as the ground colour and some markings of forewing and abdominal markings are concerned. But it differs from *flavens* Moore with respect to merged medial and postmedial bands which give almost uniform appearance and hindwing with irregular markings. As discussed under the remarks of genus the present species could not be identified from the relevant literature and by comparison with the identified collections of Natural History Museum, London. The species is unique and does not conform to the characterization of other species of genus *Thyrgorina* Walker. Hence, the species has been proposed as new to science.

**Etymology**

The species is named after its type locality i.e. Malshej Ghat in the state of Maharashtra.

*THYROGORINA EXIMIA* (SWINHOE) COMB. REV. (FIG. 4)

*Alpenus eximia* Swinhoe, 1891, *Trans. Ent.Soc.*, **1891**: 137.

*Thyrgorina eximia* (Swinhoe, 1891) (Hampson, 1894), *Fauna Br. Ind. Moths*, **2**: 14.

*Diacrisia eximia* (Swinhoe, 1891) (Hampson, 1901), *Cat. Lep. Phal.*, **3**: 301.




FIGURE 4. *Thyrgorina eximia* (Swinhoe) comb. rev. (A) Forewing; (B) Hindwing; (C) Male genitalia; (D) Aedeagus; (E) Valva (right); (F) Valva (left); (G) Uncus with Tegumen (dorsal view); (H) Uncus with Tegumen (lateral view); (I) Juxta (enlarged).



**Male genitalia**

Uncus long and narrow, slightly curved at tip, well sclerotized, setae absent; acrotergite weakly developed; fenestrula triangular; tegumen as long as uncus, u-shaped; vinculum shorter than tegumen, well sclerotized, u-shaped; saccus developed. Valvae simple, with inner walls unequally sclerotized; cucullus produced laterally to form a spine like structure. Transtilla sclerotized bar like; juxta pear shaped; aedeagus moderately narrow and long; vesica membranous, cornuti in the form of group of spines; ductus ejaculatorius entering subapically.

**Wing span** Male 30–32 mm.

**Material examined**

Karnataka: Ganeshgudi (480 m), 15.xi.03 – 1♂, 21.vii.07 – 1♂; Jogfalls (480 m), 22.vii.04 – 7♂♂.

**Old distribution** India: Uttar Kannada.

**Remarks**

The male and female genitalic structures of the species under reference have been studied and illustrated in considerable details for the first time. These structures completely conform to the characterization of genus *Thyrgorina* Walker; therefore, the old stable combination of this species has been revived in the present studies.

**Taxonomic segregation of studied taxa**

For the easy segregation of species studied here, the following key has been devised.

**KEY FOR SEPARATION OF SPECIES STUDIED**

1. Fore and hindwing with ground colour white or creamy white; abdomen with dorsal, lateral and sublateral series of black spots ..... **2**
  - Fore and hindwing with ground colour except than white; abdomen without sublateral series of black spots ..... **3**
2. Head with frons black; vertex yellow; male genitalia with juxta oval with a protrusion at distal end; female genitalia with corpus bursae funnel like .. ***indica* (Guer) comb. rev.**
  - Head with frons and vertex white; male genitalia with juxta almost rectangular with a protrusion towards distal end and a notch at proximal end; female genitalia with corpus bursae oval ..... ***angularis* (Strand) stat. nov.**
3. Head, thorax, forewing and hindwing with ground colour ochreous; abdomen orange yellow; male genitalia with uncus broad at base, slightly wavy at ventroproximal region, setosed with small setae ..... ***malshejensis* sp. nov.**
  - Head, thorax, forewing and hindwing with ground colour orange red; abdomen crimson; male genitalia with uncus comparatively narrow at base, setae absent ***eximia* (Swinhoe) comb. rev.**

## ABBREVIATIONS

IA: First anal vein; 2A: Second anal vein; AED: Aedeagus; AMP+HRP: Ampulla & Harpe (fused); ANT.APO: Anterior apophyses; CO: Costa; CRN: Cornuti; CRP.BU: Corpus Bursae; CU: Cucullus; CU<sub>1</sub>: First cubital vein; CU<sub>2</sub>: Second cubital vein; DU.BU: Ductus Bursae; Du.EJ: Ductus Ejaculatorius; FEN: Fenestrula; Jx: Juxta; M<sub>1</sub>: First median vein; M<sub>2</sub>: Second median vein; M<sub>3</sub>: Third median vein; PAPA: Papilla Analis; PO.APO: Posterior apophyses; R<sub>1</sub>: First radial vein; R<sub>2</sub>: Second radial vein; R<sub>3</sub>: Third radial vein R<sub>4</sub>: Fourth vein; R<sub>5</sub>: Fifth radial vein; RS: Radial Sector; SA: Saccus; SC: Subcosta; SC+R<sub>1</sub>: Stalk of SC + R<sub>1</sub>; SIG: Signum; SL: Sacculus; TG: Tegumen; UN: Uncus; VES: Vesica; VIN: Vinculum; VLA: Valvula; VLV: Valva.

## ACKNOWLEDGEMENTS

We are very grateful to Dr. Martin Honey, Natural History Museum, London, who rendered help for the comparison of arctiid collections with the holotypes lying there. Thanks are due to Dr. V.V. Ramamurthy, Head, Division of Entomology, Indian Agricultural Research Institute for allowing the comparison with arctiid collections. The financial help given by Department of Science & Technology, New Delhi under a project entitled 'Taxonomic revision of Indian Arctiidae (Lepidoptera)' is also duly acknowledged.

## REFERENCES

- Common, I. F. B. (1970) *Lepidoptera (moths and butterflies), in the Insect of Australia*, Melbourne University Press, Melbourne, 1-866.
- Hampson, G. F. (1894) In: *Fauna of British India, Moths, including Ceylon and Burma*, Taylor and Francis Ltd., London, **vol 2**: 1-609.
- Hampson, G. F. (1901) In: *Catalogue of the Arctiidae and Agaristidae in the Collection of the British Museum*, Taylor and Francis Ltd., London, **vol 3**: 1-690.
- Koda, N. (1988) A generic classification of the subfamily Arctiinae of Palearctic and Oriental regions based on male and female genitalia (Lepidoptera: Arctiidae) Part II. *Tyo to ga* **38(3)**: 1-79.
- Robinson, G. S. (1976) The preparation of slides of Lepidoptera genitalia with special reference to Microlepidoptera. *Entomologist's Gazette* **27**: 127-132.
- Strand, E. (1922) Arctiidae: Lithosiinae. *Lepidopterorum Catalogus* **26**: 501-825.
- Walker, F. (1864) List of the specimens of Lepidopterous insects in the collection of the British Museum. *Catalogue Lepidopetera Heterocera* **31**: 30-45.
- Zimmerman, E. C. (1903) In: *Microlepidoptera Insects of Hawaii*, University Press of Hawaii, Honolulu, **vol 9**.

(Received 6 December 2007; accepted 15 February 2008)



## Evaluation of *Bacillus thuringiensis* Berliner subsp. *kurstaki* for management of lepidopteran pests of lac insect

A. K. Jaiswal\*, A. Bhattacharya, S. Kumar and J. P. Singh

Indian Institute of Natural Resins and Gums (formerly Indian Lac Research Institute),  
Ranchi 834 010, Jharkhand, India  
Email: jaiswalak59@yahoo.co.in

**ABSTRACT:** Field trials were conducted to evaluate the commercial formulation of *Bacillus thuringiensis* Berliner subsp. *kurstaki* against two key lepidopteran predators viz., *Eublemma amabilis* Moore and *Pseudohypatopa pulverea* Meyr of lac insect on rainy season crop of *rangeeni* lac insect culture reared on the trees of *Butea monosperma* (palas). *Bt* formulations with all the concentrations tried (0.008, 0.017, 0.034, 0.05, 0.07, 0.085 per cent) significantly reduced the incidence of predators and increased the yield of broodlac over control. Two sprays of *Bt* formulation (0.05%) at an interval of one month (30 and 60 days of inoculation) significantly reduce the incidence of both the lepidopteran predators and increased the yield of broodlac significantly. However, single spray of *Bt* formulation at a concentration of 0.07% will also reduce the incidence of lepidopteran predators and increase the broodlac yield significantly. In order to get the desired efficacy of biopesticide as with chemical pesticides in suppressing the predators population, the application of *Bt* formulation at a concentration of 0.07% was found to be effective. The increase in broodlac yield amongst various treatments of biopesticide varied to the tune of 4.2–10.2 fold with single spray and 3.2–11.8 fold with double spray of *Bt* formulation.

© 2008 Association for Advancement of Entomology

**KEYWORDS:** *Eublemma amabilis*, *Pseudohypatopa pulverea*, *Bacillus thuringiensis kurstaki*, *Butea monosperma*, *Kerria lacca*, lac crop

The cultivation of lac crop is an additional source of income for many poor and economically backward families in the rural areas. Like other agricultural crops, the lac crop is also susceptible to pest attack. The lac crop is attacked by a number of predators and parasitoids but the lepidopteran predators mainly *Eublemma amabilis* Moore (Noctuidae) and *Pseudohypatopa pulverea* Meyr (Blastobasidae) are intimately associated with lac culture causing around 40% damage to lac crop (Glover, 1937; Malhotra and Katiyar, 1979). Besides several cultural practices, chemical control measures for managing these lepidopteran pests have also been evolved. Endosulfan

\*Corresponding author

(Malhotra and Katiyar, 1975, 1979), ethofenprox (Jaiswal *et al.*, 2004), dichlorvos (Mishra *et al.*, 1995, 1996) and diflubenzuron (Bhattacharya *et al.*, 1997) are some recommended insecticides which are safe to lac insect and are effective in management of the lepidopteran pests, if applied at proper time and concentration. Though the quick knock down effect have advantages over the other methods in terms of quality and quantity output specially when there is fear of rain these pesticides may pose environmental hazards in long term use. Microbial control of insect pest is one of the eco-friendly approaches without much harm to environment and *Bacillus thuringiensis* sub sp. *kurstaki* is the most popular microbial agent especially for lepidopteran pests. In order to minimize the use of chemical pesticides, *B. thuringiensis* formulation (Delfin) was evaluated on lac culture against both major lepidopteran predators.

The rainy season crop of *rangeeni* lac (June/July–Oct/Nov) was raised on trees of *Butea monosperma* (*palas*) in the month of July by inoculating broodlac. The lac insect settled on the shoots of the tree within 15 days and the used up broodlac was removed from the trees within 21 days. There were ten treatments including untreated control, each replicated thrice. Six concentrations of *B. thuringiensis* Berliner subsp. *kurstaki* (0.008, 0.017, 0.034, 0.05, 0.07 and 0.085 per cent) along with recommended concentrations of chemical pesticides viz. endosulfan 0.05%, dichlorvos 0.03% and ethofenprox 0.02% were applied. Water was sprayed in untreated control. The formulation was applied on lac culture growing on shoot of host tree with the help of a rocking sprayer. Two treatments with each concentration viz. single spray at 30 days and two sprays at 30 and 60 days of inoculation was applied. In case of dichlorvos, first spray was applied at 0.02% and second at 0.03% concentration. The randomly selected lac culture samples measuring one meter consisted of lac encrusted sticks from each tree collected at the time of crop maturity in the month of October. They were caged in 60 mesh nylon nets for quantification of incidence of insect predators. The observations on number of *E. amabilis* and *P. pulvereae* emerging from caged lac samples were recorded after 45 days of caging.

The data (Table 1) revealed that the *Bt* formulation at a concentration of 0.017 to 0.085 per cent significantly reduced the incidence of *E. amabilis* as compared to control, similar to that found with endosulfan, ethofenprox and dichlorvos. There was no significant difference between control and lowest concentration of *B. thuringiensis* (0.008 per cent). No significant difference in the reduction of predator incidence was observed with single and two sprays indicating that one spray is sufficient to minimise the incidence of *E. amabilis*. Similar results were observed with all the three insecticides. This may be due to fact that the rainy season crop is a short duration crop (110–115 days) and the mating period may be over before second spraying and the development of lac insect reaches a very advanced stage. The larvae of *E. amabilis* concealed within the lac encrustations for feeding the living lac insect and the effect of treatment is not manifested.

The incidence of *P. pulvereae* was also significantly lower than the control with all concentrations of *Bt* formulations tried as with three recommended insecticides indicating that *P. pulvereae* is more susceptible to *Bt* than *E. amabilis*. However, in

TABLE 1. Effect of *Bacillus thuringiensis* vis-à-vis other recommended pesticides on lepidopteran predators of lac insect

Insecticide	Conc. (%)	Average no. of <i>Eublemma</i> <i>ambulilis</i> per meter of lac		Average no. of <i>Pseudohypatopa</i> <i>pulverea</i> per meter of lac		Yield of brood lac (Output to input ratio) <sup>2</sup>	
		Single spray	Two sprays	Single spray	Two sprays	Single spray	Two sprays
<i>Bacillus thuringiensis</i>	0.008	7.90 (20)	7.55 (24)	4.17 (48)	5.97 (23)	1.63 (4.62)	1.50 (4.17)
	0.017	4.44 (55)	3.56 (56)	4.01 (50)	5.92 (26)	1.51 (4.21)	2.29 (6.90)
	0.034	3.83 (61)	3.48 (57)	4.00 (50)	5.33 (73)	1.64 (4.66)	2.04 (6.03)
	0.05	3.01 (70)	2.92 (64)	4.35 (46)	0	2.60 (7.97)	3.70 (11.76)
	0.07	1.85 (81)	1.00 (88)	1.62 (80)	0	2.27 (6.83)	2.75 (8.48)
	0.085	1.93 (81)	0.52 (94)	1.83 (77)	0	3.24 (10.17)	4.12 (3.21)
	0.05	1.90 (81)	0.51 (94)	3.54 (56)	0	4.00 (12.79)	2.87 (8.90)
Edosulfan	0.02	2.32 (77)	1.33 (84)	1.54 (81)	0	1.54 (4.31)	2.10 (6.24)
Ethofenprox	0.02 & 0.03	4.33 (56)	4.21 (47)	2.40 (70)	1.33 (83)	2.68 (8.24)	2.24 (6.72)
Dichlorvos		9.89	9.89	8.0	8.0	0.29	0.29
Control							
SEM ±		0.73	0.73	0.67	0.67	0.46	0.46
CD (5%)		2.07	2.07	1.90	1.90	1.31	1.31

<sup>1</sup>Figures in parentheses are % reduction over control<sup>2</sup>Fold increase in yield over control.

respect of one and two sprays, there was no significant difference in the incidence of *P. pulvereae* with lower concentrations of *Bt* ( $\geq 0.03\%$ ) but with higher concentration, the incidence differ significantly. There was no incidence of *P. pulvereae* in lac culture with two sprays of *Bt* ( $\leq 0.05\%$ ) and with endosulfan and ethofenprox. The higher sensitivity of *P. pulvereae* in comparison to *E. amabilis* may be due to the fact that the larvae of *P. pulvereae* may be better exposed to insecticides and biopesticides and also feed on dead lac insect as well as lac encrustation, thereby ingesting more spores of *Bt*. The lowest concentration which effectively reduced the incidence of *P. pulvereae* with two applications was found to be 0.05 per cent *Bt* and it is equally effective as recommended concentration of chemical pesticides namely endosulfan and ethofenprox.

The increase in broodlac yield amongst different treatments varied from 4.2–12.8 fold with single spray and 3.2–11.8 fold with double sprays in comparison to control. The yield ratio (input to output) was significantly higher than control with 0.034% and higher concentrations of *B. thuringiensis* but there was no difference between single and two sprays. There was no significant difference in yield between 0.05% endosulfan and 0.085% *Bt* treated lac crop. But yield with lower concentration of *Bt* ( $< 0.085\%$ ) was significantly less than with endosulfan. Hence in order to get the same result in respect to lac yield as with endosulfan, a 0.085% concentration of *Bt* have to be applied. There was no significant difference in yield between 0.02% ethofenprox, dichlorvos and 0.07% or less concentration of *Bt*. Hence it is inferred that *Bt* at a concentration of 0.07% and less gave same results as ethofenprox and dichlorvos. It may be concluded that in order to have better quality of broodlac (less infested with lepidopteran predators) *Bt* at a concentration of 0.05% will be suitable for double spray and 0.07% for single spray.

#### ACKNOWLEDGEMENTS

We are indebted to Department of Biotechnology, Government of India for funding the project. Encouragement by the Director, Indian Institute of Natural Resins and Gums is also acknowledged with gratitude. Thanks are also due to Shri P. Patamajhi for assistance in field and laboratory work.

#### REFERENCES

- Bhattacharya, A., Jaiswal, A. K., Sharma, K. K., Mishra, Y. D. and Chandrika, P. (1997) Evaluation of diflubenzuron on *Eublemma amabilis* (Lepidoptera: Noctuidae) a predator of lac insect *Kerria lacca* (Kerr). *Journal of Entomological Research* **21**(4): 365–369.
- Glover, P. M. (1937) *Lac Cultivation in India*, Indian Lac Research Institute, Ranchi, p. 147.
- Jaiswal, A. K., Bhattacharya, A., Kumar, K. K. and Kumar, S. (2004) Evaluation of ethofenprox (Nukil 10 EC) on lac insect culture for management of *Eublemma amabilis* Moore- a lepidopteran predator. In: *New Horizon of Animal Sciences*, Pandey, B. N., Natarajan, P., Arumugam, N. and Premjith, S. (Eds). Zoological Society of India, Magadh University, Bodh Gaya, Bihar (India), 47–50.

- Malhotra, C. P. and Katiyar, R. N. (1975) Control of *Eublemma amabilis* Moore, a serious predator of the lac insect *Kerria lacca* (Kerr) I: screening of insecticides for their safety to lac insects. *Indian Journal of Entomology* **37**: 385–396.
- Malhotra, C. P. and Katiyar, R. N. (1979) Chemical control of the lac predator *Eublemma amabilis* Moore II. Relative toxicity of the various insecticides against the predatory caterpillars. *Indian Journal of Entomology* **41**(2): 190.
- Mishra, Y. D., Bhattacharya, A., Sushil, S. N., Krishan K., Sharma and Jaiswal, A. K. (1995) Efficacy of some insecticides against *Eublemma amabilis* Moore, a major predator of lac insect, *Kerria lacca* (Kerr). *Journal of Entomological Research* **19**(4): 351–355.
- Mishra, Y. D., Sushil, S. N., Krishan K., Sharma, Bhattacharya, A. and Jaiswal, A. K. (1996) Efficacy of selected organophosphorous insecticide for control of *Chrysopa madestes* (Neuroptera: Chrysopidae)- a serious sporadic predator of Indian lac insect *Kerria lacca* (Kerr). *New Agriculturist* **7**(1): 7–20.

(Received 20 November 2007; accepted 15 February 2008)







## Effect of some selected biopesticides on growth and development of *Spodoptera Litura* fab. (Lepidoptera: Noctuidae) larvae

**M. Ravi\*, N. Dhandapani, N. Sathiah and M. Murugan**

*Department of Agricultural Entomology, Centre for Plant Protection Studies Tamil Nadu Agricultural University, Coimbatore 641003, Tamil Nadu, India*

### ABSTRACT:

Third instar larvae of *Spodoptera litura*, Fab. were fed on sunflower leaves treated with biopesticides viz., *Bacillus thuringiensis* subsp. *kurstaki* Berliner (Btk), nucleopolyhedrovirus of *S. litura* (SINPV) and azadirachtin. Food consumed was the lowest in Btk treatment. Extended feeding periods were noticed in SINPV and azadirachtin treatments. The adverse effects of biopesticides on the growth and development of *S. litura* revealed their suitability at appropriate stage in sustainable management of this pest. © 2008 Association for Advancement of Entomology

**KEYWORDS:** *Spodoptera litura*, biopesticides, food consumption, sunflower

In recent years, sunflower, *Helianthus annuus* L. is widely grown as an oilseed crop in irrigated and rainfed conditions (DES, 2004). The tobacco caterpillar, *Spodoptera litura* Fab. (Lepidoptera: Noctuidae) as a defoliator assumes the major pest status on sunflower and results in yield reduction. Management of this insect is envisaged with conventional chemical insecticides. However, use of chemical insecticides disturbs the natural balance of crop ecosystem greatly by posing problems to natural enemies and nontarget organisms. An alternative pest management technique is the use of ecofriendly biological insecticides. Larvicidal activity, antifeedant and IGR properties of microbials and botanicals are extensively proved against this pest. However, delayed mortality of the pests caused by most of the biopesticides results in continued feeding and consequent damage to crops. The present study was undertaken to find out the efficacy of selected biopesticides viz., *Bacillus thuringiensis* subsp. *kurstaki* Berliner (Btk), nucleopolyhedrovirus of *S. litura* (SINPV) and azadirachtin. Their effect on the feeding period, food consumption and utilization by the larvae were examined with a view to assess their importance in a sustainable pest management programme.

Disease free, healthy, freshly moulted third instar larvae of *S. litura* weighing 20–30 mg individually reared on semisynthetic diet (Shorey and Hale, 1965; Subrama-

\*Corresponding author

nian, 2003) were taken from the culture maintained in the Biocontrol laboratory, Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore. The larvae were released in polypots containing fresh sunflower leaves obtained from pot culture plants of known weight that were pretreated with different biopesticides viz., *Bacillus thuringiensis* subsp. *kurstaki* Berliner (Btk) (Delfin<sup>®</sup> 25 WG, Margo Biocontrols Pvt. Ltd., Bangalore), nucleopolyhedrovirus of *S. litura* (SINPV-CBE strain1) and azadirachtin (Econeem plus<sup>®</sup>, Margo Biocontrols Pvt. Ltd., Bangalore) using an atomizer and shade dried for ten minutes. Four replications were maintained per treatment with ten larvae per replication. This experiment was performed at room temperature ( $27 \pm 1^\circ\text{C}$ ) and relative humidity (70–75 %) in 12 h light, 12 h dark. Initial fresh weights of individual larva were recorded. The weight of larvae, fresh uneaten leaves and faecal matter were recorded at every 24 h interval from the first day of experiment till death/pupation of the test larvae. The weight gain or loss by the individual larva was computed at the end of the experiment. Food ingested was determined by subtracting the weight of uneaten food from weight of food provided. Collective faecal weight was also assessed. Growth indices were worked out as proposed by Waldbauer (1968).

The food consumption levels varied in the different treatments (Table 1). The consumption rate was the highest in untreated check followed by the SINPV and azadirachtin and the lowest in Btk treated food. The same trend was observed for the weight of faeces voided. The feeding period was the maximum in untreated check and minimum in the Btk treatments.

The highest consumption index (CI) was recorded for the larvae that fed on sunflower leaves treated with azadirachtin @ 500 and 1000 ppm and the lowest in Btk treatment (Table 1). The relative growth rate (RGR) was higher in azadirachtin and SINPV treatments. The efficiency of conversion of ingested food into body substances (ECI) was higher in Btk treatments followed by untreated check. Azadirachtin treatment recorded lowest ECI. The recorded net efficiency or ECD was higher in Btk treatments followed by untreated check. The AD was higher in Btk and untreated check. Azadirachtin and SINPV recorded lower AD values.

The food consumption was the lowest in Btk treated food, which might be due to the damage caused to the midgut by Btk which serves as stomach poison leading to quicker mortality (Adang, 1991). The enhanced growth rate observed with azadirachtin and SINPV on *S. litura* might be due to the development of larva in a prolonged disease suppressed condition than the normal ones (Galande and Ajri, 1997). In spite of the feeding deterrence showed by azadirachtin, the insect showed higher relative growth rate, which may be due to the altered metabolic regime in the test insect. Nucleopolyhedrovirus infected larvae have been reported to show prolonged juvenility, as the virus influenced the host via juvenile hormone activity (Ramakrishnan and Chaudhari, 1974; Subramanian, 2003).

The higher utilization of food material by *S. litura* in terms of ECI was on Btk treated food and the ECD or net efficiency was also higher in the same treatment. Naturally this condition refers to the stress situation caused by the pathogen, wherein

TABLE 1. Effect of selected biopesticides on feeding period and food consumption of third instar larvae of *Spodoptera litura*

Treatment	Feeding period (days)	Food ingested (g/larvae)	Faeces voided (g/larvae)	Consumption index (CI)	Growth rate (GR)	ECl (%)	ECD (%)	AD (%)
<i>B. thuringiensis</i> var. <i>kurstaki</i> @ 250 ppm	1.8 <sup>a</sup>	0.1564 <sup>a</sup>	0.0608 <sup>a</sup>	0.3271 <sup>d</sup>	0.0929 <sup>d</sup>	28.39 <sup>a</sup>	46.44 <sup>a</sup>	61.13 <sup>a</sup>
<i>B. thuringiensis</i> var. <i>kurstaki</i> @ 500 ppm	1.4 <sup>a</sup>	0.0986 <sup>a</sup>	0.0420 <sup>a</sup>	0.3574 <sup>d</sup>	0.0892 <sup>de</sup>	24.95 <sup>b</sup>	43.46 <sup>b</sup>	57.40 <sup>b</sup>
SINPV @ $1 \times 10^7$ POB ml <sup>-1</sup>	4.6 <sup>c</sup>	1.5294 <sup>d</sup>	0.7496 <sup>d</sup>	1.1365 <sup>b</sup>	0.1437 <sup>c</sup>	12.65 <sup>f</sup>	24.80 <sup>f</sup>	50.99 <sup>c</sup>
SINPV @ $1 \times 10^8$ POB ml <sup>-1</sup>	3.8 <sup>b</sup>	1.0304 <sup>bc</sup>	0.5246 <sup>b</sup>	0.9220 <sup>c</sup>	0.1636 <sup>b</sup>	17.74 <sup>d</sup>	32.38 <sup>d</sup>	49.09 <sup>d</sup>
Azadirachtin @ 500 ppm	4.6 <sup>c</sup>	1.2564 <sup>c</sup>	0.6488 <sup>c</sup>	1.6136 <sup>a</sup>	0.2237 <sup>a</sup>	13.87 <sup>e</sup>	28.67 <sup>e</sup>	48.36 <sup>d</sup>
Azadirachtin @ 1000 ppm	3.6 <sup>b</sup>	1.0160 <sup>b</sup>	0.5364 <sup>b</sup>	1.5979 <sup>a</sup>	0.1576 <sup>b</sup>	9.86 <sup>g</sup>	20.89 <sup>g</sup>	47.20 <sup>e</sup>
Untreated check	10.2 <sup>d</sup>	1.9810 <sup>e</sup>	0.9612 <sup>e</sup>	0.3899 <sup>d</sup>	0.0789 <sup>e</sup>	20.24 <sup>c</sup>	39.32 <sup>c</sup>	51.48 <sup>c</sup>
CD (0.01)	0.6997	0.3151	0.0595	0.0696	0.0105	1.0421	1.2793	1.0877

rates of ECD will be always faster than those of normal ones (Soo and Fraenkel, 1966). Further, it may be attributed that the cessation of feeding upon disease initiation and associated regurgitation by the larva might also have contributed to reduced food intake and feeding for a shorter period.

Index of digestibility level (AD) was highest for Btk. The reason attributed is the shortened feeding time on the Btk (Chandra *et al.*, 1999; Gujar *et al.*, 2000; Praveen and Dhandapani, 2001). Remaining treatments recorded lower AD values due to delayed mortality. It may be concluded that the *S. litura* larva is able to continue its feeding activity in spite of the contamination of the food material with biorationals indicating the slower killing effects of the biorational pesticides. However, the rate of feeding, levels of digestion and degree of assimilation varied with different biopesticides. Among them, Btk offered quicker mortality than SINPV and azadirachtin. The damage caused by surviving larvae on leaves treated with all biopesticides was significantly lower than control. Given that the damage inflicted by the intoxicated larva is tolerable, for the long term benefits and for a sustained ecological pest management, the tested biorationals stand good.

#### REFERENCES

- Adang, M. J. (1991) *Bacillus thuringiensis* insecticidal crystalproteins: Gene structure, action and utilization. In: *Biotechnology for Biological Control of Pests and Vectors*, Maramorosch, K. (Ed). CRC Press, Boca Raton, 3–24.
- Chandra, A., Kaushik, N. C. and Gupta, G. P. (1999) Studies of *Bacillus thuringiensis* on growth and development of *Helicoverpa armigera* Hubner. *Annals of Plant Protection Sciences* 7: 154–158.
- DES, (2004) *Agricultural statistics at a glance*. Directorate of Economics and Statistics (DES), New Delhi, India, 76.
- Galande, S. M. and Ajri, D. S. (1997) Consumption, digestion and utilisation of sunflower leaves by *Helicoverpa armigera* (Hubner) treated with nuclear polyhedrosis virus. *Journal of Biological Control* 11: 11–16.
- Gujar, G. T., Vinay, K. and Kumari, A. (2000) Bioactivity of *Bacillus thuringiensis* against the American bollworm, *Helicoverpa armigera* (Hubner). *Ann. Pl. Protec. Sci.* 8: 125–131.
- Praveen, P. M. and Dhandapani, N. (2001) Consumption, digestion and utilization of tomato fruits by *Helicoverpa armigera* Hubner. *Journal of Biological Control* 15: 59–62.
- Ramakrishnan, N. and Chaudhari, S. (1974) Effect of nuclear polyhedrosis disease on consumption, digestion utilization of food by the tobacco caterpillar, *Spodoptera litura* (Fab.). *Indian Journal of Entomology* 36: 101–105.
- Shorey, H. H. and Hale, R. L. (1965) Mass rearing of the larvae of nine noctuid species on a simple artificial medium. *Journal of Economic Entomology* 58: 522–524.
- Soo, H. C. F. and Fraenkel, G. (1966) The consumption, digestion and utilisation of food plants by a polyphagous insect, *Prodenia eridania* (Cramer). *Journal of Insect Physiology* 12: 711–730.
- Subramanian, S. (2003) Studies on the use of baculoviruses for the management of *Plutella xylostella* (L.) and *Spodoptera litura* (F.) infesting cauliflower. Unpub. *Ph.D. Thesis*, Tamil Nadu Agricultural University, Coimbatore, India, p 152p.
- Waldbauer, G. P. (1968) The consumption and utilization of food by insects. *Advances in Insect Physiology* 5: 229–288.

(Received 4 May 2006; accepted 15 February 2008)



## Occurrence of Phthiraptera on the house crow, *Corvus splendens* (Passeriformes: Corvidae)

Sultan Beg<sup>1</sup>, Nidhi Gupta<sup>1</sup>, Sandeep Kumar<sup>2</sup>, Vikram Khan<sup>1</sup>,  
Shivika Bhatnagar<sup>1</sup> and A K Saxena\*<sup>1</sup>

<sup>1</sup>Department of Zoology, Govt. Raza P.G. College, Rampur, Uttar Pradesh, India

<sup>2</sup>Department of Zoology, Govt. P.G. College, Gopeshwar, Dist. Chamoli, Uttarakhand, India

Email: akscsir@rediffmail.com

**ABSTRACT:** The prevalence and other population parameters of five phthirapteran species on a sample of 70 house crows in the district Rampur, U.P., India were determined. The most prevalent species was *Philopterus lahorensis* (51.45% of the population), followed by *Allocolpocephalum fregili* (34.3%), *Corvonirmus saliem* (30.0%), *Menacanthus gonophaeus* (18.6%) and *Myrsidea baktitar* (11.4%). All the five species exhibited aggregated distribution but the distribution did not conform to negative binomial model. Females outnumbered the males in all the five species; the adult to nymph ratio varied among the species.

© 2008 Association for Advancement of Entomology

**KEYWORDS:** Phthiraptera, Mallophaga, biting lice, population parameters on crow

Prevalence and frequency distribution pattern of Phthiraptera on selected avian hosts has been recorded from time to time. Rozsa *et al.* (1996) compared the louse loads of territorial hooded crows (*Corvus coronae cornix*) and colonial rooks (*C. frugilegus*). A survey of literature indicated that there was no report on the population levels of Phthiraptera infesting the house crows (*Corvus splendens*). The present report furnishes information on the prevalence and frequency distribution patterns of five phthirapteran species occurring on house crow.

For the present studies, fumigation method described by Clayton and Drown (2001) was adopted. As a deviation, polythene bags and chloroform were used instead of Anaesthesia jar and ethyl acetate, suggested in the method. The entire louse load was transferred to 70% alcohol and separated according to species, stage and sex, for further analysis. The data were used for determining the prevalence, mean intensity, sample mean abundance and variance to mean ratio of the louse population. The exponent (k) of negative binomial distribution was estimated with the help of software

\*Corresponding author

offered by Rozsa *et al.* (2000)). The goodness of fit between the observed and the expected frequencies (negative binomial) was determined by chi-square test.

A total of 2345 specimens belonging to five species viz. *Allocolpocephalum fregili* Denny, *Menacanthus gonophaeus* Burmeister, *Myrsidea baktitar* Ansari, *Corvonirmus* (= *Brueelia*) *saliemi* Ansari and *Phlopterus lahorensis* Ansari were recovered. Price *et al.* (2003) recognized only five phthirapteran species on the aforesaid host (*B. saliemi*, *C. fregili*, *P. lahorensis*, *Menacanthus gonophaeus* and *Myrsidea merisui* Eichler). In the present studies, *M. merisui* was not detected although *M. baktitar* was recovered. Price *et al.* (2003) have mentioned the presence of *M. baktitar* on a subspecies of *C. splendens* (*C. s. zugmayeri*).

Prevalence of Phthiraptera on house crows was 67.5% ( $n = 70$ ). The overall mean intensity of Phthiraptera on the house crows remained 49.9 while the sample mean abundance was 33.5. *P. lahorensis* was the most prevalent species. The prevalence of different species in the order of descending frequency was *P. lahorensis* > *A. fregili* > *C. saliemi* > *M. gonophaeus* > *M. baktitar* (table 1). The mean intensities and sample mean abundances varied from 10.6 to 27.0 and 1.2 to 13.9, respectively. The values of variance to mean ratio exceeded unity in the case of all the five species. The prevalence of different phthirapteran species on Hungarian rooks and hooded crows (Rozsa *et al.*, 1996) appear to differ from that of house crows, recorded during present study. Furthermore, the infestation intensity of different species on Hungarian rooks (1–15) and hooded crows (3.9–4.9) were comparatively lower than house crows (10.6–27.0).

The values of negative binomial exponent ( $k$ ) ranged from 0.03 to 0.2. The frequency distribution patterns were aggregated in case of all the five species but the observed frequencies failed to correspond the frequencies expected by negative binomial model. Survey of literature indicated that phthirapteran ectoparasites generally exhibit aggregated (clumped) distribution as most of the hosts have few parasites while a few have lots of them. However, Rekasi *et al.* (1997) found that distribution patterns of twenty one (out of 27 species, occurring on 13 birds) conformed to negative binomial model. But Saxena *et al.* (2007) and Gupta *et al.* (2007) reported that the frequency distribution pattern of only one species (out of 12 species, occurring on five avian hosts) conformed with the negative binomial model.

An examination of population composition (table 1) shows that female population outnumbered the males in case of all the five species. However, the adult to nymph ratios were quite inconsistent. It remained nearly equal in case of *A. fregili* and *C. saliemi*. Nymphal population dominated over adults in case of *P. lahorensis*. However, in case of *M. gonophaeus*, the adults outnumbered the nymphs in natural population. The examination of population composition of any species provides useful information about the temporal stability of the population; there will be fewer adults and more nymphs if the population is increasing (Marshall, 1981). However, the population composition (proportion of juveniles) is bound to vary with time, due to several environmental and host factors (Marshall, 1981). As far as sex ratio is concerned, there was considerable uniformity as the females outnumbered the males in case of all the

TABLE 1. Population of five species of lice on house crows (*Corvus splendens*) in district Rampur, U.P., India

Observations	<i>Allocoipocephalum fregili</i>	<i>Menacanthus gonophaes</i>	<i>Myrsidea baktitar</i>	<i>Philopterus lahorensis</i>	<i>Corvonirmus sallemi</i>
Total number of lice recovered	556	259	85	971	474
Total number of males	122	53	17	164	90
Total number of females	151	84	33	222	139
Total number of nymphs	283	122	35	585	245
Adult nymph ratio	1:1	1:0.9	1:0.7	1:1.5	1:1.1
Male female ratio	1:1.2	1:1.6	1:1.9	1:1.4	1:1.5
Ratio of nymphal stages	1:0.9:0.7	1:0.8:0.7	1:0.8:0.6	1:0.9:0.7	1:0.9:0.8
Number of birds infested	24	13	8	36	21
Prevalence (%)	34.3	18.6	11.4	51.4	30
Mean Intensity	23.2	19.9	10.6	27.0	22.6
Sample mean abundance	7.9	3.7	1.2	13.9	6.8
Variance/ Mean ratio	30.3	29.5	16.1	25.0	23.7
Range of infestation	5-72	1-58	2-26	2-56	3-48
k of negative binomial	0.11	0.05	0.03	0.2	0.09
Negative binomial (Yes/ No)	No	No	No	No	No

species. In phthirapteran population, sex ratios are generally female biased (Marshall, 1981). Unequal longevity (males generally short lived than females), sampling errors (being small sized, males are easily lost during sampling) and local mate competition (if female lice disperse more effectively than males) presumably cause imbalances in sex ratios of phthirapteran population (Marshall, 1981; Rozsa *et al.*, 1996; Saxena *et al.*, 2007).

#### ACKNOWLEDGEMENTS

We are thankful to the Principal, Govt. Raza P. G. College, Rampur for laboratory facilities and to the Department of Science & Technology, New Delhi, India for providing financial support to A.K.S. in the form of project No. SP/SO/AS-30/2002

#### REFERENCES

- Clayton, D. H. and Drown, D. M. (2001) Critical evaluation of five methods for quantifying chewing lice (Insecta: Phthiraptera). *Journal of Parasitology* **87**(6): 1291–1300.
- Gupta, N., Kumar, S. and Saxena, A. K. (2007) Prevalence and population structure of lice (Phthiraptera) on the Indian Red Avadavat. *Zoological Science* **24**: 381–383.
- Marshall, A. G. (1981) *The Ecology of Ectoparasitic Insects*, Academic Press, London, p. 459.
- Price, R. D., Hellenenthal, R. A., Palma, R. L., Johnson, K. P. and Clayton, D. H. (2003) The chewing lice: World checklist and biological overview. In: *Illinois Natural History Survey Special Publication 24*, Champaign, Illinois, p. 501.
- Rekasi, J., Rozsa, L. and Kiss, B. J. (1997) Patterns in the distribution of avian lice (Phthiraptera: Amblycera, Ischnocera). *Journal of Avian Biology* **28**: 150–156.
- Rozsa, L., Rekasi, J. and Reiczigel, J. (1996) Relationship of host coloniality to the population ecology of avian lice (Insecta: Phthiraptera). *Journal of Animal Ecology* **65**: 242–248.
- Rozsa, L., Reiczigel, J. and Majoros, G. (2000) Quantifying parasites in samples of hosts. *Journal of Parasitology* **86**: 228–232.
- Saxena, A. K., Kumar, S., Gupta, N., Mitra, J. D., Ali, S.A. and Srivastava, R. (2007) Distribution pattern of phthirapterans infesting certain common Indian birds. *Journal of Parasitology* **93**(4): 957–958.

(Received 8 October 2007; accepted 15 February 2008)





## Effect of row spacing and different levels of nitrogenous fertilizer on population density of thrips (*Thrips tabaci* Lind.) in onion (*Allium cepa* L.)

Abhishek Shukla<sup>\*†</sup> and S. S. Rathore

Krishi Vigyan Kendra, Chomu 303702, Jaipur, Rajasthan, India  
Email: isdp.udaipur@sify.com

**ABSTRACT:** A field experiment was conducted to find out the effect of row spacing and different levels of nitrogenous fertilizers on the population densities of thrips (*T. tabaci*) infesting onion. A spacing of 30 cm between rows and 150 kg/ha nitrogen had low level of thrips population and gave highest yield.

© 2008 Association for Advancement of Entomology

**KEYWORDS:** thrips, row spacing, fertilizers, population densities

Onion (*Allium cepa* L.) is an important condimental bulbous crop. It is a major horticultural cash crop grown in Rajasthan. Thrips (*Thrips tabaci* Lind.) is a major pest of this crop which causes considerable yield loss. Various agronomic and cultural practices play an important role in population build up of pest, slight modification in these practices will some times reduce the population density of insect pests greatly. Keeping the point trials were conducted to investigate the effect of different row spacing and different levels of nitrogenous fertilizers on the population density of thrips (*T. tabaci*) in onion crop.

The experiments were laid out at Experimental Farm of Krishi Vigyan Kendra, Chomu, Jaipur, Rajasthan during *rabi* 2005–06 and 2006–07 in randomized block design (RBD). Onion variety RO-59 was used for the trials.

An experiment was conducted to evaluate row to row spacing effect on thrips population in onion for two consecutive years, with following five treatments of row spacing i.e., 20, 25, 30, 35 and 40 cm. respectively. The seedlings were transplanted in 24 equal plots (8 × 10 m). All the treatments were free from insecticidal application. Agronomic practices were followed as per the recommendation. Data on the thrips population were recorded weekly from five randomly selected plants from each plot

\*Corresponding author

†Present address: Fruit Research Station, Navsari Agricultural Univeristy, Gandevi 396360, Gujarat, India.

TABLE 1. Effect of row spacing on population densities of thrips in onion

Treatment (Row spacing in cm)	No. of thrips per plant		Yield (q/ha)	
	2005-06	2006-07	2005-06	2006-07
20 × 20	8.290	8.453	191.225	179.575
25 × 25	6.945	6.650	255.103	234.985
30 × 30	6.100	6.373	265.875	240.815
35 × 35	4.805	4.605	214.088	218.643
40 × 40	4.598	4.515	211.688	214.200
CD at 5%	0.236	0.182	56.222	73.432

TABLE 2. Effect of different levels of nitrogenous fertilizers on population densities of thrips in onion

Treatments (N:P:K)	No. of thrips per plant		Yield (q/ha)	
	2005-06	2006-07	2005-06	2006-07
(0:100:100)	7.618	7.820	108.225	100.650
(50:100:100)	7.518	7.455	120.988	123.025
(100:100:100)	7.588	7.668	141.675	135.625
(150:100:100)	7.855	7.928	155.613	144.625
(200:100:100)	13.293	14.238	111.325	111.525
(250:100:100)	13.343	13.793	94.933	89.213
CD at 5%	0.218	0.438	2.250	2.115

by plant wash method described by Bullock (1963) until the crop was harvested. After digging, yield from each plot was recorded.

A separate trial was conducted to evaluate the influence of five different levels of nitrogenous fertilizers on thrips population. The trial was conducted with six treatments — one control without NPK and remaining by the varying recommended dose of NPK fertilizers. N levels as 50, 100, 150, 200 and 250, half applied at planting and balance 8 weeks after planting with P and K given full as basal dose. For counting thrips the plant wash method described by Bullock (1963) was followed. Yield from each plot was also recorded.

The maximum number of thrips (8.290 and 8.453 per plant) was observed in 20 × 20 cm spacing and the minimum in 35 × 20 cm and 40 × 20 cm spacing. The maximum yield (265.875 and 240.815 q/ha) came from 30 × 30 cm spacing. The lowest yield was recorded in 20 × 20 cm spacing during both the years. Edelson *et al.* (1986) reported a negative relationship between thrips population and onion yield. Yield data did not significantly vary in both the years.

Table 2 shows the influence of different rates of N with constant P, K levels on thrips population densities during the years 2005-06 and 2006-07. No effect was observed on thrips densities up to 150 kg/ha, while 200 and 250 kg/ha had the maximum

thrips densities (13.93 and 14.238; 13.343 and 13.793 thrips per plant, respectively). Fertilizers provide plants with more nutrients (Bentz *et al.*, 1995) as a result of which the plants not only get a lush green colour but also enhance the accumulation of nutrients in the shoots, which attract phytophagous insects. Use of fertilizers not affects the nutritive value of plants but also impacts the insect pests densities (Bentz and Larew, 1992). Onion yield showed a direct relation to the increased fertilizer rates up to 150 kg/ha. The maximum fertilizer rate of 250 kg/ha gave minimum yield of 89.213 q/ha. Excessive use of chemical fertilizers might have caused disturbance in physiological activities of the plants that adversely affect the yield (Megdi *et al.*, 2001). The treatment 150 kg/ha gave the maximum yield (155.613 and 144.625 q/ha) during both the years of study. For the proper growth of plants an optimum rate of nutrients is very important (Nashrin *et al.*, 2001). On the basis of the results it is clear that thrips are not a threat if the fertilizers are used in low or optimal rates. Under the present study the 150–100–100 NPK kg/ha is the best fertilizer rate for onion yield and low thrips infestation.

#### REFERENCES

- Bentz, J. A. and Larew, H. G. (1992) Oviposition preference and nymphal performa of *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae) on *Dendranthema grandiflora* under different fertilizer regimes. *Journal of Economic Entomology* **85**: 514–517.
- Bentz, J. A., Reeves, J., Barbosa, P. and Francis, B. (1995) Nitrogen fertilizer effect on selection, acceptance and suitability of *Euphorbia pulcherrima* (Euphorbiaceae) as a host plant to *Bemisia tabaci* (Homoptera: Aleyrodidae). *Environmental Entomology* **24**: 40–45.
- Bullock, J. A. (1963) Extraction of Thysanoptera from samples of foliage. *Journal of Economic Entomology* **56**: 612–614.
- Edelson, J. V., Cartwright, B. and Royer, T. A. (1986) Distribution and impact of *Thrips tabaci* (Thysanoptera: Thripidae) on onion. *Journal of Economic Entomology* **79**: 502–505.
- Megdi, S. M., Shakir, A., Sadiq, M., Sarfaraz, M., Hassan, G., Akhtar, J. and Jamil, M. (2001) Effect of phosphorus, zinc and farm yard manure in the presence of nitrogen and potash on NP and Zn concentration in rice. *Pakistan Journal of Biological Science* **4**: 342–343.
- Nashrin, S. A., Farooqi, A. M., Siddiqui, M. A., Rahman, M. M. and Khanam, M. R. (2001) Effect of nitrogen and spacing on the growth and yield of *Gimakalmi ipomoea* Reptans Poir. *Journal of Biological Science* **2**: 170–174.

(Received 13 August 2007; accepted 15 February 2008)





## Pests of economic importance on *Jatropha curcas* L., a biodiesel plant in Andhra Pradesh, India

**M. Prabhakar\*, Y. G. Prasad, G. R. Rao and D. Ramakrishna**

*Central Research Institute for Dryland Agriculture, Hyderabad 500059, Andhra Pradesh, India*

*Email: mprabhakar@crida.ernet.in*

**ABSTRACT:** Large-scale cultivation of *Jatropha curcas* is being taken up in Andhra Pradesh, India as a source of biodiesel. An attempt for an inventory of pests causing significant damage to *J. curcas* plant is made for the first time in Andhra Pradesh. Damage by the twig girdler, *Oncideres limpida* on *J. curcas* is reported for the first time in India. Five species of insects viz., thrips, *Retithrips syriacus* (Mayet); spotted plant bugs, *Scutellera nobilis* (Fabricius) and *Chrysocoris purpureus* (Westwood); soft scale, *Megapulvinaria maxima* (Green); inflorescence webber and capsule borer, *Pempelia morosalis* (Saalm Uller) were also seen severely infesting *Jatropha curcas* (L.) in Andhra Pradesh. Their seasonal incidence, nature and extent of damage also have been described. © 2008 Association for Advancement of Entomology

**KEYWORDS:** *Jatropha curcas*, biodiesel, pests, Andhra Pradesh

*Jatropha curcas* (L.), commonly known as physic nut is a medicinal plant, hedge crop and also widely used in soap industry. More importantly, the oil extracted from seeds is a promising substitute for diesel. As per the estimates of Ministry of Rural Development, Government of India the potential area for *Jatropha* cultivation in Andhra Pradesh is 43.96 lakh ha, and it is the target of Government of Andhra Pradesh to raise biofuels in 2 lakh ha across the state by 2008–09 (Kureel, 2006). Information available on the pest scenario on *J. curcas* from Andhra Pradesh is inadequate for the future crop protection needs relating to the crop. Hence, the present study was aimed at making an inventory of pests that cause economic loss to *J. curcas* in Andhra Pradesh. Fortnightly observations were made during 2005–06 on 30 randomly selected plants from a 3 ha block of two year old *J. curcas* in the Hayathnagar mandal of Rangareddy district, Andhra Pradesh. No pesticides were sprayed on the selected plants.

*Oncideres limpida* (Coleoptera: Cerambycidae), the twig girder was one of the serious pests observed on *J. curcas*. Branches close to ground at two feet height and having thickness between 12 and 21 mm were the worst affected. The adult beetle chews a continuous notch girdling the twig. The girdles are similar to that made by

\*Corresponding author

a saw and the ends appear gnawed almost straight across with a faint rounding. The girdled twig soon dries out and breaks off; and the larva develops inside the fallen twig on the ground. The incidence of girdler was recorded mainly during October–November. The number of affected branches on each plant was in the range of 2–8 and about eight percent of the plants in the experimental block were found affected. The female beetles of *O. limpida* damaging *Prosopis juliflora* has been reported earlier from Rajasthan (Singh, 1993). This is the first report of *O. limpida* on *J. curcas* in India.

*Retithrips syriacus* (Mayet) (Thysanoptera: Thripidae) occur as colonies on leaves and tender fruits. Adults and nymphs cause damage by scraping the epidermal tissue and suck sap from leaves and fruits, resulting in white patches, which are visible from a distance. The damaged portions on the fruits turn into black patches and often lead to secondary infection of powdery mildew. Severe foliar infestation resulted in premature leaf drop. It occurs all through the year except March and April. Peak incidence was in May–June and again in November–December. The mean number was in the range of 35 to 250 and adult nymph ratio was 3.2: 29.4. Heavy rains during July–August diminished their population and the plants recovered their natural vigour. This species was first reported in India by Seshadri and Ananthakrishnan (1954). Later it was reported on many crops viz., grapevine, groundnut, pomegranate, and cashew and rose (Nair, 1975; Dash and Naik, 1998). However, the incidence of *R. syriacus* on *Jatropha* was first reported by Manoharan *et al.* (2006) from Tamil Nadu and the present report is for the first time from Andhra Pradesh.

*Scutellera nobilis* (Fabricius) and *Chrysocoris purpureus* (Westwood) (Scutellariidae: Hemiptera) are the two species of bugs found infesting *J. curcas*. Both the bugs are metallic blue-green in colour with 8–12 black spots on the dorsal side of thorax and abdomen. Their incidence was noticed throughout the year, but high population was recorded during August–November and the mean number of adults per plant was between 2 and 15. Eggs are pink in colour, found in groups on the leaves. On hatching the pale red grubs with black patches on thorax move all around the plant in groups and suck sap from leaves, flowers and fruits. Feeding on inflorescence caused premature flower drop. Attack on fruits left pinholes, which further lead to rotting of the rind and eventually fruit drop. The seeds from affected fruits were chaffy. *S. nobilis* has been reported infesting tamarind in Madhya Pradesh (Meshram and Garg, 1999) and cotton in Haryana (Sucheta and Khokhar, 2004). On *J. curcas* the incidence of *S. nobilis* had been reported from Tamil Nadu (Manoharan *et al.*, 2006) and Uttar Pradesh (Chitra Shankar and Dhyani, 2006) in India.

Infestation by soft scales *Megapulvinaria maxima* (Green) (Hemiptera: Coccidae) causing white scabs on leaves and twigs were common. All stages except the newly hatched nymphs are sessile. Movement of ants on the plants was the first indication of scale infestation. *M. maxima* incidence was observed from August to February. The peak infestation was in October and November. Infested plants often showed growth of black sooty mold. Severe incidence of *Pulvinaria maxima* on *Jatropha* sp. and other plants like neem, cotton and mulberry was reported by Ayyar (1919) in

Andhra Pradesh. It was reported as a serious pest on neem (Annamalai *et al.*, 1996) and more recently on *J. curcas* in one location out of the three surveyed in Tamil Nadu (Manoharan *et al.*, 2006).

The larvae of inflorescence webber and capsule borer, *Pempelia morosalis* (Lepidoptera: Pyralidae) occur in groups and feed on leaves, inflorescence and tender buds by making silken webs. The damage was severe on inflorescence and fruits. The seeds in the affected fruits were either partly or completely damaged resulting in heavy yield loss. *Jatropha* is an indeterminate crop and it comes to flowering many times a year depending on climate and availability of soil moisture. Peak incidence of webber was recorded in February–March and August–September, coinciding with flowering on *Jatropha*. The number of webs per plant ranged between 3 and 11 and the number of larvae in each web between 6 and 30. *Salberia (Pempelia) morosalis* (Saalm Uller) was first reported from India as leaf and flower webber on few forest plants including *Jatropha* by Beeson (1941). More recently it has been reported as capsule borer and bark-eating caterpillar on *J. curcas* from Uttar Pradesh (Chitra Shankar and Dhyani, 2006). This is the first report of *P. morosalis* infesting *J. curcas* in Andhra Pradesh.

#### ACKNOWLEDGEMENTS

We are grateful to NOVOD Board, Gurgaon for financial assistance and the Director and Heads DCS and DRM, CRIDA, Hyderabad for providing necessary facilities. The services rendered by Prof. C. A. Viraktamath, UAS, Bangalore and Prof. S. Suresh, TNAU, Coimbatore in identifying the insect specimens is gratefully acknowledged.

#### REFERENCES

- Annamalai, R., Surendran, C., Thirumurthi, S. and Gunasekharan, V. (1996) Insect damage to neem in Tamil Nadu, India. In: *Impact of diseases and insect pests in tropical forests. In: Proceedings of IUFRO Symposium, Peechi, India, 23-26 November, 1993*, 343–355.
- Ayyar, T. V. R. (1919) A contribution to our knowledge to south Indian Coccidae. *Bulletin of the Agricultural Research Institute, Pusa* **87**: 1–50.
- Beeson, C. F. C. (1941) *The Ecology and Control of the Forest Insects of India and Neighbouring Countries*, Forest Research Institute, Dehra Dun, India, p. 1007.
- Chitra Shankar, S. K. and Dhyani, (2006) Insect pests of *Jatropha curcas* L. and the potential for their management. *Current Science* **19(2)**: 162–163.
- Dash, P. C. and Naik, A. K. (1998) Thrips fauna associated with rose in Orissa. *Insect Environment* **4(2)**: 59.
- Kureel, R. S. (2006) Biofuel scenario in India. In: *In Proceedings of the Third International Biofuels Conference, 18–19 January 2006*, New Delhi, India, 20–51.
- Manoharan, T., Ambika, S., Natarajan, N. and Senguttuvan, K. (2006) Emerging pest status of *Jatropha curcas* (L) in south India. *Indian Journal of Agroforestry* **8**: 66–79.
- Meshram, P. B. and Garg, V. K. (1999) A report on the occurrence of *Scutellera nobilis* Fab. on *Emblica officinalis* Gaertn. *Indian Forester* **125(5)**: 536.
- Nair, M. R. G. K. (1975) *Insects and Mites of Crops in India*, ICAR, New Delhi, India, 273 pp.
- Seshadri, A. and Ananthakrishnan, T. N. (1954) Some new Indian Thysanoptera - I. *Indian Journal of Entomology* **16**: 210–226.

- Singh, M. P. (1993) Injurious hexapoda associated with prosopis. In: *Proceedings of a Conference on Prosopis species in the Arid and Semi-Arid Zones of India*, Central Arid Zone Research Institute, Jodhpur, Rajasthan, India, 21–23.(November 1993)
- Sucheta, K. and Khokhar, K. S. (2004) Biosystematics and biometric studies of pentatomid bugs (Hemiptera: Pentatomidae) associated with cotton. *Journal of Cotton Research and Development* **18**(2): 215–219.

(Received 26 December 2007; accepted 15 February 2008)





## **Incidence of shot hole borer, *Xylosandrus compactus* Eichhoff (Coleoptera: Scolytidae) on wilt disease affected cocoa**

**K. Subaharan<sup>1</sup>, C. Kannan<sup>2</sup>, Rohini Iyer<sup>1</sup> and P. S. P. Vidyasagar<sup>1</sup>**

<sup>1</sup>*Central Plantation Crops Research Institute, Kasaragod 671 124, India*

<sup>2</sup>*Central Plantation Crops Research Institute (RS) Vittal, Karnataka, India*

**ABSTRACT:** Extensive wilting of cocoa trees has been reported from Mysore, Chamrajanagar, Bangalore and Mandya districts of Karnataka. The wilted trees showed shot hole borings made by *Xylosandrus compactus* on the dead twigs. All the stages of the pest were found in the galleries made by the pest in twigs. The life cycle was completed in 28 days in laboratory. Twigs of 11–20 mm thickness were preferred over those below 10 mm and above 30 mm in thickness. Since the incidence was observed only on dead twigs the association of pest incidence and the wilting could not be established. © 2008 Association for Advancement of Entomology

**KEYWORDS:** *Xylosandrus compactus*, cocoa wilt, *Ceratocystis* sp.

Wilting and rapid death of cocoa (*Theobroma cocoa*) trees has been reported from Mysore, Mandya, Chamrajanagar and Bangalore districts of Karnataka during 2000 (Chowdappa *et al.*, 2002). The etiology of this disease is still unknown. Occurrence of shot holes caused by *Xylosandrus compactus* was observed in the field in Mysore district of Karnataka. In India there is no record of this species infesting cocoa but it has been recorded as a primary pest on coffee seedlings (Muthappa and Venkatasubbaiah, 1981). Hence preliminary studies were carried out and the findings are reported here.

The occurrence of beetle infestation in relation to stem thickness was assessed. Twenty twigs of 25 cm length and girth of 5–10 mm, 10–20 mm, 20–30 mm and > 30 mm were collected randomly from 20 plants in wilt infested gardens. Collection was made randomly from all the four directions. Mean number of shot holes in these samples was recorded in relation to the girth of the stem. The mean holes in twigs having 5 to 10, 11 to 20, 28 to 30 and > 30 mm thickness were 0.8, 4.05, 2 and 0 respectively.

The twigs having the bore holes contained all stages of the pest in the gallery. The pest was reared in the laboratory on wilted cocoa twigs to study the biology. The egg,

larval and pupal periods were observed as 3.8, 5.3 and 8.66 days, respectively, the total being 28.5 days.

In India *X. compactus* was reported as a primary pest on robusta coffee and its attack was confined to seedlings (Muthappa and Venkatasubbaiah, 1981). In cocoa the incidence was more on 4–5 years old trees. The biology of this pest was reported on coffee in Ivory Coast (Brader, 1962) and Hawaii (Hara and Beardsley, 1979) and on cocoa in Nigeria (Gregory, 1954). Sixty species of ambrosia beetles have been recorded to infest cocoa twigs and majority of them have been recorded as secondary feeders. Of these only *X. ferrugineus* F. attacks healthy cocoa. *X. corniculatus* Schedl and *X. posticus* Eich. attack diseased or dead trees only (Entwistle, 1972).

In Karnataka the incidence of *X. compactus* was restricted to wilt affected trees. Hara and Beardsley (1979) reported that stressed plants were more susceptible to infestation by scolytids. The attraction of ambrosia beetles (especially *Xyleborus*) to diseased plants of *Mangifera* and *Eucalyptus* were reported earlier (Entwistle, 1972). Association if any between *Xylosandrus compactus* and the wilt disease of cocoa in Karnataka can be established only with further research.

#### ACKNOWLEDGEMENTS


The authors are thankful to Cadburys India for assisting in field observations at Yelwala. The technical assistance provided by Mr. P. Ravindran, CPCRI, Kasaragod is acknowledged.

#### REFERENCES

- Brader, L. (1962) Observations on life history of *Xyleborus morstatti* in the Ivory Coast. *Tijdscher. Pl Zeiekt* **69**: 111–113.
- Chowdappa, Rohini, Iyer and Vidyasagar, P. S. P. V. (2002) Cocoa wilt a major threat to cocoa cultivation in traditional areas of India. In: *Abst. of Natl Symp. Crop Prot. and WTO*, 22–25, January, 2002, CPCRI, Kasaragod.
- Entwistle, P.F. (1972) *Pests of Cocoa*, Longman, UK.
- Gregory, J.L. (1954) Nigeria. Review of the work by the Entomological section, Dept. of Agriculture, 1948 – 54. *Rep. 6th Commonw. Ent. Conf.* 295–296.
- Hara, A.H. and Beardsley, J.W. Jr (1979) The biology of the black twig borer, *Xylosandrus* (Eichhoff), in Hawaii. *Pro. Hawaiian Entomol. Soc.* **18**(1): 55–70.
- Muthappa, B.N. and Venkatasubbaiah, P. (1981) Association of *Ambrosiella macrospora* with *Xylosandrus compactus*, the shot-hole borer of robusta coffee in India. *J. Coffee Res.* **11**(2): 54.

(Received 27 November 2007; accepted 15 February 2008)

## REVIEW OF BOOK



## **Tropical forest insect pests**

**Ecology, Impact, and Management**

**K. S. S. Nair**

Cambridge University Press, UK

Published 10 May 2007, 422 pp.

Though we have a few recent books published in Tropical Forest Entomology, the present one under review is unique because it is written by an author, who has a first hand field knowledge on the various issues related to forest pest problems in the tropics. Moreover, Dr. K. S. S. Nair with his abilities both as a teacher and researcher is an apt person to write a book like this.

The forest is an abode of rich biodiversity of insects. With the onset of large scale forest plantations coming up, it has caused pest problems as well, which are different from that of the agriculture sector. Thus developing pest control strategies in forest plantations is a challenging problem. In forestry, considerable data on the pest ecology would be required to decide on even the methods to be adopted for managing the pest. Thus success stories on pest management in forestry are limited.

As the author himself has indicated in the text “Forest entomology is rich in theories”. It has to be further added that most of these are based on case studies from temperate forests. In contrast, the examples which are lucidly presented in this book are from the tropics.

The introductory pages of this book set the stage for discussing the varied interactions between the forest insects and their habitat. The initial chapter describes the tropics, tropical forest types, species diversity, forest growth dynamics, management practices, non-timber forest products and also issues such as deforestation, sustainable management, plantations and exotics. Under structural diversity, 30 orders are

presented with details on six major insect orders of forestry importance. Seven feeding guilds are also presented to explain the functional diversity. The term pest has also been adequately explained. The functional role of insects in the flow of matter and energy in the forest ecosystem is described and also the two existing theories on insect pest outbreaks- 'Resource supply theory' and 'Host concentration theory' are explained.

The book also has a separate chapter on pest incidence in natural forests with many case studies from across the globe. The insect pest problems in tropical forest plantations have also been covered in detail. There is a very useful chapter on the pest scenario of twenty four tree species giving the details of the tree species, pest spectrum, impact of pests, management aspects and knowledge gaps. This chapter will be an updated source of reference for researchers, teachers and students of forest entomology.

Three much debated and controversial aspects – pest incidence in natural forests vs. plantations, exotics vs. indigenous tree species and monocultures vs. mixed plantations, have been dealt with in a very balanced manner.

Most of the entomologists have a pet experimental insect and in forestry, choosing a pet insect is quite difficult. But Dr. Nair, because of his inherent interest in population dynamics of insects, rightly chose the teak defoliator, *Hyblaea puera* as his pet insect and worked relentlessly towards finding out a viable control strategy by understanding the ecology of this pest. The formulation of the pest evolution hypothesis to explain the greater incidence of pest outbreaks in plantations than in natural forests is helpful in evolving control strategies. *H. puera* is a major pest problem in most of the teak growing countries, especially in the tropics. The descriptions given on various aspects of the ecology of this pest shows his in depth field knowledge and they are exactly like what you find in a field situation.

Illustrations and photographs in the book are excellent and would have been more impressive, if printed in colour. The book also contains over 500 references of great relevance for anyone to read and understand more on specific topics in forest entomology. Thus this book is an important addition to the tropical forest entomology literature which would enthuse youngsters to take up serious studies on the ecology and management of forest insect pests.

Excerpts from the book can be viewed at [www.cambridge.org/9780521873321](http://www.cambridge.org/9780521873321)

Dr. R.V. Varma  
Member Secretary  
Kerala State Biodiversity Board  
Pallimukku, Pettah  
Thiruvananthapuram 695 024

## GUIDELINES TO AUTHORS

**Scope:** ENTOMON will publish original research papers on insects, arachnids and myriapods. Reviews are not acceptable.

Papers on morphology, anatomy and histology will be considered only when they form part of systematics, physiology or behavioural studies.

Announcements of seminars/symposia, book reviews and other items of entomological interest will also be considered for publication.

**Types of papers:** Articles up to 3 printed pages in length (1800 words) will be published as **short communications** and those between 4 and 10 pages (up to 6000 words) as **full papers**.

Full paper should be organized under the following sub-heads—Abstract, Introduction, Materials and Methods, Results, Discussion, Acknowledgements and References.

Short communication should be organized in the same way, but without the sub-headings.

**Publication policy:** Manuscripts submitted for publication in ENTOMON should not have been published or submitted for publication elsewhere.

At least one of the authors should be a member of AAE.

Page charges are applicable at the rate of Rs. 100 per printed page for authors in India and US \$ 10 for authors overseas. Invoice will be sent to the authors along with the galley proof.

Twenty five **reprints** of the paper will be supplied free of cost to the authors.

**Typing and submission:** Manuscripts should be typed double space on one side of good quality paper, having 3.5 cm left margin and 2.5 cm right margin.

The first page should contain the title, author's name affiliation and e-mail address. When the number of authors is more than one, indicate the name of the corresponding author with an asterisk and specify 'Corresponding author' in a footnote. The second page should contain the abstract, followed by key words and a running title. From page 3 onwards, type the text continuously from Introduction to References. Place the tables on separate sheets at the end of the paper. The pages should be numbered.

Three copies of the manuscript, complete in all respects including illustrations, should be sent to the Managing Editor, ENTOMON, Department of Zoology, University of Kerala, Kariavattom P.O., Thiruvananthapuram 695 581, Kerala, India.

An electronic version need be sent only after acceptance of the paper for publication.

**Guide for writing:** Careful attention to the following guide will facilitate early acceptance of your paper for publication in ENTOMON. Although some of these suggestions may appear trivial, they have been prompted by our experience in reviewing the papers received for publication. Keep in mind that ENTOMON is a research journal and your paper will be read only by those who are specialists in the respective fields of research.

**Title** should be brief and should reflect the specific content of the research reported in the paper.

**Abstract** should be informative, not indicative. It should very briefly highlight the aim of the study and major conclusions. Normally it should not exceed 150 words.

**Key words** should be limited to four or five most pertinent indicators of the work, relevant to indexing the article.

**Introduction** should be brief, normally not exceeding 300 words. Include the specific aim of the research, a review of the available information in the area and existing gap in knowledge. The introduction should thus justify the work carried out. Avoid elementary details and repetition of well known facts. For example, in a paper reporting the efficacy of a biopesticide against a particular pest, it is not necessary to explain the hazards of chemical pesticides, alternative methods of insect control and advantage of integrated pest management. Limit the literature review to what is most relevant to the topic under study.

**Materials and Methods** should provide just enough details to permit proper interpretation of the results. Materials used need not be described separately if it is evident from the methods given. Technical description of the method is needed only when the method is new. If the method followed has been already described elsewhere, just give the reference. If any alteration is made, describe the alteration alone, with reason.

**Results:** Adequate care should be taken to organize and present the results in a clear, concise and summarized form.

Quantitative data should always be analysed using suitable statistical methods. Organize the data into well planned tables. Each table should be self-explanatory.

Do not repeat the data presented in the table in the text. Quote the relevant figures in the text only when it is essential for highlighting some particular finding.

Due care should be taken while interpreting the results of statistical analysis. For example, treatments which show higher numerical value cannot be treated as superior to those having lower numerical values when there is no statistically significant difference.

Interpretation of the data should be with reference to the objectives set in the experiment.

Do not include graphs duplicating the data presented in the tables.

When the research involves repetition of the work already reported by others, include the new findings alone in the paper.

**Illustrations** included in the paper should be essential for explaining some points in the text. Photographs of the life stages of an insect are not useful unless the insect is being reported for the first time. Illustration should be of good quality. Limit the number of photographs to 4–6 and organize them into plates wherever possible. The illustrations should be numbered consecutively as Fig. 1, Fig. 2, etc., without distinction between drawings, graphs and photographs. Labelling should be legible and large enough to stand suitable reduction. Legend for the figures should be typed on a separate page. All figures must be referred to, at appropriate places, in the text.

The cost of printing colour illustration is to be met by the author.

**Discussion:** The discussion section is intended to critically analyse and interpret the results with reference to the objectives set forth in the study. It should highlight the importance of the results in relation to what is already known. It should also point out the limitations of the study, if any. The discussion should not repeat details given under Results, except to highlight some conclusions.

**References** should list all publications cited in the text, in alphabetical order. Do not include any references not cited in the text. Some authors show a tendency to cite too many references in support of a statement. Cite only a few references most pertinent to the point dealt with. Follow the citation style given below.

*Examples of citations in text:*

Krishnaswamy (1978, 1979)

Govindan *et al.* (1998)

(Reddy 1978; David 1991)

*Examples of citations under References:*

Articles in Journals:

Nayar K. K. (1953) Neurosecretion in *Iphita*. *Current Science* 22(2): 149.

Nair M. R. G. K. and Mohandas N. (1962) On the biology and control of *Carvalhoeia arecae*, a pest of areca palms in Kerala. *Indian Journal of Entomology* 24: (1) 86–93.

Jalaja M. Muraleedharan D. and Prabhu V. K. K. (1973) Effect of extirpation of median neurosecretory cells on reproduction in the female red cotton bug, *Dysdercus cingulatus*. *Journal of Insect Physiology* 19(1): 29–36

Books and Articles in Books:

Novak V. J. A. (1966) *Insect Hormones*. Methuen and Co., 478 pp.

Wigglesworth V. B. (1964) The hormonal regulation of growth and reproduction in insects. In: *Advances in Insect Physiology* Vol. 2 (Eds. Beament J. W. L., Treherne J. E. and Wigglesworth V. B), Academic Press, London, pp 247–335.

**Review Procedure:** Manuscripts will be subjected to a preliminary scrutiny by an Editorial team and those conforming to the general guidelines set above will be referred to experts for detailed review; others will be returned to the author for resubmission in the prescribed format. Based on detailed review, the corresponding author will be advised of acceptance, non-acceptance or need for revision. The revised

manuscript will be scrutinized again by an Editorial team (and by expert referees, if needed) before final acceptance. On final acceptance, the author will be asked to submit an electronic version of the manuscript. Proof will be sent to the corresponding author. It should be checked and returned within three days of receipt. The journal reserves the right to proceed with publication if corrections are not communicated promptly.

**Strict conformity with the above guidelines will  
ensure speedy publication of your paper.**



## AUTHOR INDEX

Padmavathi, I	Leena, S., 41
Anandhi, 47	Mohan, Murali, 25
Beg, Sultan, 75	Muraleedharan, D., 41
Bhatnagar, Shivika, 75	Murugan, M., 71
Bhattacharya, A., 65	Padmakumari, A. P., 1
Dhandapani, N., 71	M I. C., 1
Gill, Navneet Singh, 53	Pillai, M. A. K., 47
Gupta, Nidhi, 75	Prabhakar, M., 83
Iyer, Rohini, 87	Prasad, Y. G., 83
Jacob, Sosamma, 35	Ramakrishna, D., 83
Jaiswal, A. K., 65	Rao, G. R., 83
Joshi, Sunil, 15	Rathore, S. S., 79
Kannan, C., 87	Ravi, M., 71
Karthikeyan, K., 35	Rayudu, B. T., 41
Katti, Gururaj, 1	Sathiah, N., 71
Khan, Vikram, 75	Saxena, A K, 75
Kirti, Jagbir Singh, 53	Shukla, Abhishek, 79
Kumar, S., 65	Singh, J. P., 65
Kumar, Sandeep, 75	Subaharan, K., 87
	Subaharan, K., 25
	Varma, Savita, 47
	Vidyasagar, P. S. P., 87

---

### Statement of ownership and other particulars about ENTOMON

(Form IV, Rule 8 of Registration of Newspapers (Central) Rules 1956)

1. Place of publication:	Trivandrum
2. Periodicity of publication:	Quarterly
3. Printer's name, nationality: and address:	D. Muraleedharan, Indian CABB, University of Kerala Kariavattom, Trivandrum 695581
4. Publisher's name, nationality and address:	-do-
5. Editor's name, nationality and address:	-do-
6. Name and address of the individual who owns the newspaper:	Association for Advancement of Entomology CABB, University of Kerala Kariavattom, Trivandrum 695581

I, D. Muraleedharan, hereby declare that the particulars given above are true to the best of my knowledge and belief.

Trivandrum  
31 March 2008

Sd/-  
D. Muraleedharan  
Publisher, ENTOMON

Pests of economic importance on <i>Jatropha curcas</i> L., a biodiesel plant in Andhra Pradesh, India: M. Prabhakar, Y. G. Prasad, G. R. Rao, D. Ramakrishna. . . . .	83
Incidence of shot hole borer, <i>Xylosandrus compactus</i> Eichhoff (Coleoptera: Scolytidae) on wilt disease affected cocoa: K. Subaharan, C. Kannan, Rohini Iyer, P. S. P. Vidyasagar. . . . .	87